arXiv:1001.4670v2 [math.GR] 5 Dec 2011

ON VOLUMES OF ARITHMETIC QUOTIENTS OF $\mathrm{PO}(n,1)^\circ$, n ODD

MIKHAIL BELOLIPETSKY AND VINCENT EMERY

ABSTRACT. We determine the minimal volume of arithmetic hyperbolic orientable n-dimensional orbifolds (compact and non-compact) for every odd dimension $n \geq 5$. Combined with the previously known results it solves the minimal volume problem for arithmetic hyperbolic n-orbifolds in all dimensions.

1. INTRODUCTION

In this paper we study volumes of arithmetic hyperbolic orbifolds of odd dimension. Our main results determine the arithmetic hyperbolic orientable n-dimensional orbifolds (compact and non-compact) of smallest volume for every odd dimension $n \geq 5$. This work can be considered as a continuation of the research which was done in [6] for $n = 3$ and in [2] for even $n > 2$. The question of minimal volume for $n = 2$ is answered by a classical theorem of Siegel [28]. Together with these results our paper concludes the study of the minimal volume for arithmetic hyperbolic n-orbifolds in all dimensions. Our Theorems 1 and 2 give a more precise version of the results of the second author's PhD thesis [9]. We refer to the thesis for an introduction to the methods used in this paper and for a more detailed exposition of the proofs.

Let $\mathbb{H}^n$ be the hyperbolic n-space and let $\mathrm{Isom}(\mathbb{H}^n)$ be its group of isometries. Given a lattice Γ in $\mathrm{PO}(n,1) = \mathrm{Isom}(\mathbb{H}^n)$, the corresponding quotient space $\mathbb{H}^n/\Gamma$ is called a *hyperbolic n-orbifold*. When $\Gamma < \mathrm{Isom}^+(\mathbb{H}^n) = \mathrm{PO}(n,1)^\circ$, the orbifold is orientable. If Γ is an arithmetic lattice defined over a number field k (see Section 2.2), we call $O = \mathbb{H}^n/\Gamma$ an *arithmetic orbifold* and we call k its *field of definition*.

Theorem 1. *For each dimension $n = 2r-1 \geq 5$, there is a unique orientable compact arithmetic hyperbolic n-orbifold O_0^n of the smallest volume. It is defined over $k_0 = \mathbb{Q}[\sqrt{5}]$ and has hyperbolic volume given by the formula*

$$\mathrm{Vol}_{\mathbb{H}}(O_0^n) = \frac{5^{r^2-r/2}\cdot 11^{r-1/2}\cdot (r-1)!}{2^{2r-1}\pi^r}\, L_{\ell_0|k_0}(r) \prod_{i=1}^{r-1} \frac{(2i-1)!^2}{(2\pi)^{4i}}\,\zeta_{k_0}(2i),$$

where ℓ_0 is the quartic field with a defining polynomial $x^4 - x^3 + 2x - 1$.

2010 *Mathematics Subject Classification.* 22E40 (primary); 11E57, 20G30, 51M25 (secondary).
Belolipetsky partially supported by EPSRC grant EP/F022662/1.
Emery partially supported by SNSF, project no. 200020-121506/1 and fellowship no. PBFRP2-128067.

The proof of Theorem 1 is conceptually similar to that in [2]: The main ingredients are Prasad's volume formula [25], some results and ideas of Borel-Prasad [5] and Bruhat-Tits theory (see [31]). Essential differences in comparison with the previous work lie on the technical side. They can be explained by considering the algebraic group whose real points is $\mathrm{PO}(n,1)^\circ$. First, its algebraic simply connected covering is a 4-covering when n is odd (while it is only a double covering when n is even). Secondly, when n is odd, $\mathrm{PO}(n,1)^\circ$ is of type D, for which there exist outer forms (for n even the type is B and all forms of this type are inner). In particular, the triality forms ${}^{3,6}\mathrm{D}_4$ come out while dealing with $n = 7$. However, our work shows that the orbifold O_0^7 is not defined by a triality form. In order to deal with this issues we need to analyze in detail the structure of the algebraic groups of type D defined over number fields. More precisely, compact hyperbolic arithmetic orbifolds can be constructed using any totally real number field different from $\mathbb{Q}$.

Geometric methods can be used in small dimensions ($n \leq 9$) to determine non-compact hyperbolic n-orbifolds of the smallest volume (including non-arithmetic orbifolds) [18, 13, 14]. All these minimal orbifolds are known to be arithmetic. The following theorem is consistent with the results of [13] and gives new results in higher dimensions:

Theorem 2. *For each dimension $n = 2r-1 \geq 5$, there is a unique orientable non-compact arithmetic hyperbolic n-orbifold O_1^n of the smallest volume. Its volume is given by:*

(1) if $r \equiv 1 \pmod 4$:

$$\mathrm{Vol}_{\mathbb{H}}(O_1^n) \quad = \quad \frac{1}{2^{r-2}}\,\zeta(r)\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}}\zeta(2i);$$

(2) if $r \equiv 3 \pmod 4$:

$$\mathrm{Vol}_{\mathbb{H}}(O_1^n) \quad = \quad \frac{(2^r-1)(2^{r-1}-1)}{3\cdot 2^{r-1}}\,\zeta(r)\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}}\zeta(2i);$$

(3) if r is even:

$$\mathrm{Vol}_{\mathbb{H}}(O_1^n) \quad = \quad \frac{3^{r-1/2}}{2^{r-1}}\,L_{\ell_1|\mathbb{Q}}(r)\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}}\zeta(2i),$$

where $\ell_1 = \mathbb{Q}[\sqrt{-3}]$.

The remarks made above about the proof of Theorem 1 remain valid for Theorem 2, except some specific issues. Thus, by Godement's compactness criterion the field of definition of a non-compact arithmetic hyperbolic n-orbifold is known to be the field of rational numbers $\mathbb{Q}$. Also, the issue of triality forms does not appear here: non-compact hyperbolic orbifolds are never related to the triality (see Section 2.3). Regardless of this few simplifications the proof of Theorem 2 requires essentially all the machinery which is used for Theorem 1.

The results of Hild in [13] which we mentioned above allow us to compare the formulas from Theorem 2 with what was obtained by geometric methods for small dimensions. We note that the results of [13] are not limited to orientable orbifolds. In particular, it is proved there that the smallest volume n-orbifold for $n = 5,7,9$ is non-orientable and unique. This shows that the orientable double covers of these orbifolds correspond to our O_1^n ($n = 5,7,9$). Their volumes coincide with the formulas given in Theorem 2.

By evaluating the volumes in Theorems 1 and 2 and comparing their asymptotic growth for large n we obtain the following theorem of an independent interest. It was already known for even dimensions (see [2, Sec. 4.5]), and our work allows us to extend it for all sufficiently large n. This result gives further support to a general conjecture stated in [2, Add. 1.6].

Theorem 3. *For every dimension $n \geq 5$, the minimal volume of a non-compact arithmetic hyperbolic n-orbifold is smaller than the volume of any compact arithmetic hyperbolic n-orbifold. Moreover, the ratio between the minimal volumes* $\mathrm{Vol}_{\mathbb{H}}(O_0^n)/\mathrm{Vol}_{\mathbb{H}}(O_1^n)$ *grows super-exponentially with n.*

It is interesting to point out a counterintuitive feature of this result: certain orbifolds which are non-compact and thus have infinite cusps appear to be smaller than any compact arithmetic orbifold of the same dimension. We can recall that for $2 \leq n \leq 4$ the situation is different, here the minimal volume of a non-compact hyperbolic n-orbifold is larger than the volume of the smallest compact arithmetic n-orbifold (see [6], [18], [2] and [14]). The same result holds true for $n = 2$ and 3 without arithmeticity assumption. We conjecture that in higher dimensions one can also remove the assumption of arithmeticity and that our Theorem 3 should apply to all hyperbolic n-orbifolds.

We now give a brief outline of the paper. Section 2 provides some preliminary results which are used along the lines. In Section 3 we construct principal arithmetic subgroups Λ_0 and Λ_1 whose normalizers appear to be the natural candidates for the arithmetic subgroups of minimal covolume. This construction is motivated by Prasad's volume formula introduced in Section 2.6. In Section 4 we give a bound for the index of a principal arithmetic subgroup in its normalizer. This bound is essential for our argument, its derivation is based on a detailed study of Galois cohomology. The results of Section 4 are used in Section 5 to compute the precise indices for our groups Λ_0 and Λ_1. Uniqueness of the groups Λ_0, Λ_1 and their normalizers is established in Section 6. The proof is based on a reformulation of the problem in terms of Galois cohomology which allows us to apply the methods of the class field theory together with the results of the previous sections to show the uniqueness. After all these preparations we are ready to prove the main theorems. This is done in Sections 7–10. The idea is first to use the general estimates and reduce the list of the possible candidates for the minimal volume. Then we improve the estimates using some specific properties of the candidates from the list and reiterate the procedure. After several steps of this optimization process we reduce the number of possibilities to a few candidates. A careful analysis of these remaining cases shows that the minimum is always attained on the groups which were constructed in Section 3. This finishes the proof of Theorems 1 and 2. The final Section 10 is dedicated to the study of growth of the minimal volume and the proof of Theorem 3. The results obtained here have some interesting applications. In particular, the method of [1] allows to apply these results together with [2] to the problem of classification of arithmetic hyperbolic reflection groups.

Acknowledgements. We would like to thank Ruth Kellerhals, Gopal Prasad, Alireza Salehi Golsefidy and Thomas Weigel for their interest in this work and helpful discussions. We would also like to thank the referee for their comments and suggestions.

2. PRELIMINARIES ON VOLUMES AND ARITHMETIC SUBGROUPS

2.1. The group of isometries of the hyperbolic n-space $\mathrm{Isom}(\mathbb{H}^n)$ is isomorphic to the real Lie group $\mathrm{PO}(n,1)$. The group of orientation preserving isometries $\mathrm{Isom}^+(\mathbb{H}^n)$ is isomorphic to the identity component $\overline{H} = \mathrm{PO}(n,1)^\circ$, which can be further identified with the matrix group $\mathrm{SO}(n,1)^\circ$. The simply connected covering of $\overline{H}$ is the spinor group $H = \mathrm{Spin}(n,1)$. Any orientable hyperbolic n-orbifold is a quotient $\mathbb{H}^n/\overline{\Gamma}$ for some lattice $\overline{\Gamma}$ in $\overline{H}$. We can consider the full inverse image Γ of $\overline{\Gamma}$ in H. A lattice in H can be identified with such an inverse image whenever it contains the center of H.

Let $\mathfrak{h}$ be the Lie algebra of H (and $\overline{H}$). Each Haar measure on H or $\overline{H}$ corresponds to a multilinear form of the highest degree on $\mathfrak{h}$. If K denotes a maximal compact subgroup of $\overline{H}$ with Lie algebra $\mathfrak{k}$, then $\mathbb{H}^n$ can be identified with $\overline{H}/K$. Furthermore, the tangent space of $\mathbb{H}^n = \overline{H}/K$ at the point K can be identified with a specially chosen subspace $\mathfrak{p} \subset \mathfrak{h}$ such that $\mathfrak{h} = \mathfrak{k} \oplus \mathfrak{p}$. It is known [12, Ch. V, §2] that $\mathfrak{k} \oplus i\mathfrak{p}$ is the (real) Lie algebra of the groups $\mathrm{Spin}(n+1)$ and $\mathrm{SO}(n+1)$, and this gives a duality between $\mathbb{H}^n$ and the n-sphere of curvature one. Thus for any Haar measure ν on H there is a corresponding Haar measure on $\mathrm{Spin}(n+1)$, which we will denote by ν^*. Moreover, we denote by the same symbol two Haar measures on H and $\overline{H}$ that correspond to a same multilinear form on $\mathfrak{h}$. We adopt the same convention for the measures on $\mathrm{Spin}(n,1)$ and $\mathrm{SO}(n+1)$.

Let ω^1 be the multilinear form on $\mathfrak{p}$ corresponding to the left-invariant measure $\mathrm{Vol}_{\mathbb{H}}$ and let ω^0 be the form on $\mathfrak{k}$ normalized by $\int_K \omega^0 = 1$. Then the Haar measure $\mu_{\mathbb{H}}$ given by $\omega^0 \wedge \omega^1$ is such that for any lattice $\overline{\Gamma}$ in $\overline{H}$ we have:

$$\mathrm{Vol}_{\mathbb{H}}(\mathbb{H}^n/\overline{\Gamma}) \quad = \quad \mu_{\mathbb{H}}(\overline{H}/\overline{\Gamma}).$$

Note that if Γ contains the center of H, we have $\mu_{\mathbb{H}}(H/\Gamma) = \mu_{\mathbb{H}}(\overline{H}/\overline{\Gamma})$. By the duality we can also check that $\mu_{\mathbb{H}}^*(\mathrm{SO}(n+1))$ is equal to the usual volume of the unit n-sphere.

In this article μ stands for the Haar measure on H normalized by $\mu^*(\mathrm{Spin}(n+1)) = 1$. This is the normalization of the measure used by Prasad in [25]. Since the covering $\mathrm{Spin}(n+1) \to \mathrm{SO}(n+1)$ is of degree two, we have $\mu^*(\mathrm{SO}(n+1)) = 1/2$. Hence using the formula for the volume of the unit n-sphere in $\mathbb{R}^{n+1}$, in the case of n odd, we get:

$$\mu_{\mathbb{H}} \quad = \quad \frac{4\pi^{\frac{n+1}{2}}}{\left(\frac{n-1}{2}\right)!}\,\mu. \tag{1}$$

This allows us to work with μ instead of $\mathrm{Vol}_{\mathbb{H}}$.

2.2. Let G be an absolute simple, simply connected algebraic group defined over a number field k. We denote by V_f (resp. V_∞) the set of finite (resp. infinite) places of k. A collection $\mathscr{P} = (\mathrm{P}_v)_{v \in V_f}$ of parahoric subgroups $\mathrm{P}_v \subset \mathrm{G}(k_v)$ is called *coherent* if the product $\prod_{v\in V_f} \mathrm{P}_v$ is an open subgroup of the finite adèle group $\mathrm{G}(\mathbb{A}_f(k))$ (we refer to e.g. [25, Sec. 0.5] for the definition of parahoric subgroups). In this case the intersection $\Lambda_{\mathscr{P}} = \mathrm{G}(k) \cap \prod_{v \in V_f} \mathrm{P}_v$ is called a *principal arithmetic subgroup* of $\mathrm{G}(k)$. Any subgroup of $\mathrm{G}(k \otimes_{\mathbb{Q}} \mathbb{R})$ that is commensurable with some principal arithmetic subgroup is called an *arithmetic subgroup*.

By the Bruhat-Tits theory, the conjugacy class (referred to as the *type*) of a parahoric subgroup P_v in $\mathrm{G}(k_v)$ corresponds to a subset θ_v of a basis Δ_v of the affine root system of $\mathrm{G}(k_v)$. This way a principal arithmetic subgroup Λ determines a *global type*

$$\theta = \prod_{v \in V_f} \theta_v \subset \prod_{v \in V_f} \Delta_v.$$

An algebraic group G as above will be called *admissible* if there exists a continuous surjective homomorphism $\phi : \mathrm{G}(k \otimes_{\mathbb{Q}} \mathbb{R}) \to H$ whose kernel is compact. This implies that G is simply connected and there exists $v_0 \in V_\infty$ such that $\mathrm{G}(k_{v_0}) \simeq H$ and $\mathrm{G}(k_v)$ is compact for all $v \in V_\infty \setminus \{v_0\}$. In particular, k has to be a totally real field. Given an arithmetic subgroup $\Gamma \subset \mathrm{G}(k \otimes_{\mathbb{Q}} \mathbb{R})$ for some admissible group G defined over k, its image $\phi(\Gamma)$ is a lattice in H. All subgroups of H which are commensurable with such $\phi(\Gamma)$ are called *arithmetic subgroups of H defined over k*. Arithmetic subgroups of H of the form $\phi(\Lambda)$ for some principal arithmetic subgroup Λ of G will be called the *principal arithmetic subgroups of H*.

From now on G will always be an admissible group defined over k and of absolute rank $r \geq 3$, $v_0 \in V_\infty$ denotes the archimedean place at which G is isotropic, and Δ_v $(v \in V_f)$ denotes a chosen basis of the affine root system of $\mathrm{G}(k_v)$.

2.3. The group G is of absolute type D_r (where $n = 2r-1$) and we will often write D_{2m} or D_{2m+1} to distinguish between r even or odd. It follows from the classification of algebraic groups [30] and Godement's compactness criterion that G gives rise to cocompact lattices in H if and only if $k \neq \mathbb{Q}$ (see [16, Secs. 1-2] for details).

By the admissibility condition, $\mathrm{G}(k \otimes_{\mathbb{Q}} \mathbb{R})$ contains exactly one factor $\mathrm{Spin}(n,1)$ and the rest of the factors, if any, are isomorphic to $\mathrm{Spin}(n+1)$. By the classification of semisimple algebraic groups (cf. [30, §3]) the group $\mathrm{Spin}(n,1)$ (resp. $\mathrm{Spin}(n+1)$) is an inner form exactly when the discriminant of the standard quadratic form of signature $(n,1)$ (resp. the standard positive definite form) has the same sign as the discriminant of the split form in $n+1$ variables, which equals $(-1)^r$. For even $r = 2m$ we conclude that $\mathrm{Spin}(n,1)$ is outer. This forces G to be an outer form (of possible types ${}^2\mathrm{D}_{2m}$ or the exceptional *triality forms* ${}^{3,6}\mathrm{D}_4$). For odd $r = 2m+1$ and $k \neq \mathbb{Q}$ the group $\mathrm{G}(k \otimes_{\mathbb{Q}} \mathbb{R})$ must contain at least one compact factor $\mathrm{Spin}(n+1)$, which is then an outer form. In this case again G is an outer form (of the only possible type ${}^2\mathrm{D}_{2m+1}$). For the remaining case $k = \mathbb{Q}$ with odd rank, both types ${}^1\mathrm{D}_{2m+1}$ and ${}^2\mathrm{D}_{2m+1}$ are possible.

2.4. A certain field extension $\ell|k$ attached to G will be used throughout the paper. For G of type different from ${}^6\mathrm{D}_4$, the field ℓ is defined as the minimal extension of k such that G becomes an inner form over ℓ. In particular, except for the type ${}^3\mathrm{D}_4$, we have $[\ell : k] \leq 2$. For G of type ${}^6\mathrm{D}_4$ we let ℓ to be a fixed extension of k of degree 3 which is contained in the minimal extension of k of degree 6 over which G is inner. For the types ${}^{3,6}\mathrm{D}_4$ we have $[\ell : k] = 3$. By abuse of language the field ℓ will be called in this paper the *splitting field* of G.

Let (s_1, s_2) be the *signature* of the field ℓ, i.e. s_1 (resp. s_2) is the number of real (resp. complex) places of ℓ. In the following proposition we collect some useful information about this field.

Proposition 2.5. *Let* $\mathrm{G}|k$ *be an admissible group of the absolute rank* r *and let* $d = [k : \mathbb{Q}]$. *We suppose that* G *is an outer form.*

1. *If* $r = 2m+1$, *then the signature of* ℓ *is given by* $(s_1, s_2) = (2, d-1)$.
2. *If* $r = 2m$ *and* G *is not of type* ${}^{3,6}\mathrm{D}_4$ *, then* $(s_1, s_2) = (2d-2, 1)$.
3. *When* $k \neq \mathbb{Q}$ *the field* ℓ *has at least one real place and the only roots of unity in* ℓ *are* ± 1.

Proof. The field ℓ, except for the case ${}^6\mathrm{D}_4$, is the minimal splitting field of the quasi-split inner form of G. This implies that for $v \in V_\infty$, if G is an inner form over k_v, the field ℓ is a subfield of k_v and hence for any place w of ℓ extending v the field ℓ_w is real. If G is an outer form over k_v the opposite holds: there must exist $w|v$ for which ℓ_w is complex. The two first assertions now follow from the description of the type of $\mathrm{Spin}(n,1)$ and $\mathrm{Spin}(n+1)$ given in Section 2.3.

If $k \neq \mathbb{Q}$ and the type is different from ${}^{3,6}\mathrm{D}_4$, the previous consideration proves also that ℓ must have at least one real place. For the cases ${}^{3,6}\mathrm{D}_4$ we have $[\ell : k] = 3$ and then for each $v \in V_\infty$ we must have at least one real place $w|v$ of ℓ, since $[\ell : k] = \sum_{w|v} [\ell_w : k_v]$. Thus in each of the cases ℓ has at least one real place, and hence the only roots of unity in ℓ are ± 1. This proves the third assertion. □

2.6. The covolume of a principal arithmetic subgroup $\Lambda \subset \mathrm{G}(k_{v_0}) \simeq H$ corresponding to a coherent collection $\mathscr{P} = (\mathrm{P}_v)_{v \in V_f}$ can be computed using Prasad's volume formula [25, Theorem 3.7]. In our case Prasad's formula gives

$$\mu(H/\Lambda) = \mathscr{D}_k^{\frac{2r^2-r}{2}} \left(\frac{\mathscr{D}_\ell}{\mathscr{D}_k^{[\ell:k]}} \right)^{\frac{1}{2}(2r-1)} C(r)^{[k:\mathbb{Q}]} \, \mathscr{E}(\mathscr{P}), \tag{2}$$

where ℓ is the field introduced in Section 2.4, $\mathscr{D}_K$ denotes the absolute value of the discriminant of a number field K (in this paper we will call $\mathscr{D}_K$ briefly the *discriminant* of K), $\mathscr{E}(\mathscr{P}) = \prod_{v \in V_f} e_v(\mathrm{P}_v)$ is an Euler product of local factors which are determined by the structure of $(\mathrm{P}_v)_{v\in V_f}$, and the constant $C(r)$ is defined by

$$C(r)=\frac{(r-1)!}{(2\pi)^r}\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}}.$$

2.7. The local factors $e(\mathrm{P}_v)$ can be computed using the Bruhat-Tits theory. An extensive account of its main concepts and results is given in [31].

Let k_v be a nonarchimedean local field, $\hat{k}_v$ its maximal unramified extension, and P_v be a parahoric subgroup of $\mathrm{G}(k_v)$. Following [25, Sec. 3.7], $e(\mathrm{P}_v)$ is given by the formula

$$e(\mathrm{P}_v)=\frac{q_v^{(\dim\overline{\mathrm{M}}_v+\dim\overline{\mathscr{M}}_v)/2}}{\#\overline{\mathrm{M}}_v(\mathfrak{f}_v)},$$

where q_v is the degree of the residue field $\mathfrak{f}_v$ of k_v, and $\overline{\mathrm{M}}_v$, $\overline{\mathscr{M}}_v$ are certain connected reductive $\mathfrak{f}_v$-groups associated to G and its quasi-split inner form which are defined in [25, Sec. 2.2]. There exists a minimal finite (possibly empty) subset $T\subset V_f$, such that for all $v\in V_f\setminus T$, the group G is quasi-split over k_v and P_v is a hyperspecial (resp. special) parahoric subgroup if G splits over $\hat{k}_v$ (resp. does not split over $\hat{k}_v$). For $v\in V_f\setminus T$ the group $\overline{\mathrm{M}}_v$ is isomorphic to $\overline{\mathscr{M}}_v$ over $\mathfrak{f}_v$.

We assume till the end of Section 2.7 that G is not of type ${}^{3,6}\mathrm{D}_4$. If G splits over $\hat{k}_v$ and $v\in V_f\setminus T$, then $\overline{\mathrm{M}}_v\simeq\overline{\mathscr{M}}_v$ is a connected semisimple $\mathfrak{f}_v$-group of type ${}^1\mathrm{D}_r$ or ${}^2\mathrm{D}_r$ according with G being split or non-split over k_v. If G does not split over $\hat{k}_v$ and $v\in V_f\setminus T$, then $\overline{\mathrm{M}}_v\simeq\overline{\mathscr{M}}_v$ is of type B_{r-1}. Hence for $v\in V_f\setminus T$, we have (see [23, Table 1]):

$$\dim\overline{\mathrm{M}}_v=\dim\overline{\mathscr{M}}_v=2r^2-r,$$

$$\#\overline{\mathrm{M}}_v(\mathfrak{f}_v)=q^{r(r-1)}(q^r\mp 1)\prod_{i=1}^{r-1}(q^{2i}-1),$$

if G splits over $\hat{k}_v$, the sign $\mp$ being negative (resp. positive) if $\mathrm{G}|k_v$ is inner (resp. outer); and

$$\dim\overline{\mathrm{M}}_v=\dim\overline{\mathscr{M}}_v=2(r-1)^2+r-1,$$

$$\#\overline{\mathrm{M}}_v(\mathfrak{f}_v)=q^{(r-1)^2}\prod_{i=1}^{r-1}(q^{2i}-1),$$

if G does not split over $\hat{k}_v$.

We can bring these cases together using the field ℓ defined in Section 2.4. Namely, assuming for a moment that $T=\emptyset$, we obtain

$$\prod_{v\in V_f}e(\mathrm{P}_v)=\zeta_k(2)\zeta_k(4)\ldots\zeta_k(2r-2)L_{\ell|k}(r).$$

Now, for each $v \in T$ we define a so-called *lambda factor*:

$$\lambda_v \quad = \quad e(\mathrm{P}_v) \cdot \frac{\#\overline{\mathscr{M}}_v(\mathfrak{f}_v)}{q_v^{\dim(\overline{\mathscr{M}}_v)}}. \tag{3}$$

Then

$$\mathscr{E}(\mathscr{P}) = \zeta_k(2)\zeta_k(4)\ldots\zeta_k(2r-2)L_{\ell|k}(r)\prod_{v\in T}\lambda_v, \tag{4}$$

moreover, by [25, Prop. 2.10],

$$\begin{aligned} \lambda_v &\geq (q_v+1)^{-1}q_v^{r_v+1}(1\mp q_v^{-r_v})\prod_{i=1}^{r_v-1}(1-q_v^{-2i}) \\ &\geq \frac{2}{3}\left(\frac{3}{4}q_v\right)^{r_v}, \end{aligned} \tag{5}$$

where r_v is the $\hat{k}_v$-rank of G.

2.8. It is well known that every maximal arithmetic subgroup Γ of $H = \mathrm{G}(k_{v_0})$ is a normalizer $N_H(\Lambda)$ of some principal arithmetic subgroup Λ (see e.g. [5, Prop. 1.4]). In [26] (see also [7]) Rohlfs gave a characterization of principal arithmetic subgroups whose normalizers are maximal by defining a notion of "$\mathscr{O}$-maximality". Arithmetic subgroups of minimal covolume are necessarily maximal. In order to determine these subgroups we will first look at the principal arithmetic subgroups which respect Rohlfs' criterion and have small covolume (Section 3), and later on will consider their normalizers.

3. CONSTRUCTION OF Λ_0 AND Λ_1

3.1. In this section we give a construction of the principal arithmetic subgroups Λ_0 and Λ_1, whose normalizers define the minimal orbifolds from Theorems 1 and 2. These will be subgroups of admissible spinor groups Spin_f for some quadratic forms f. The group $\mathrm{G} = \mathrm{Spin}_f$ needs to fulfill the admissibility condition of Section 2.2. This is precisely the case when f is defined over a totally real number field k, has signature $(n,1)$ at an infinite place v_0 of k and is anisotropic for all $v \in V_\infty \setminus \{v_0\}$. We call such f an *admissible quadratic form*. Arithmetic subgroups of such groups G are sometimes referred to as arithmetic lattices of the *first type*. Such a lattice is cocompact if and only if $k \neq \mathbb{Q}$ (see Section 2.3). All non-cocompact arithmetic subgroups are of the first type [16, Secs. 1-2], i.e. every admissible $\mathbb{Q}$-group is of the form Spin_f for some admissible quadratic form f defined over $\mathbb{Q}$.

3.2. Let $(\mathbf{V}, f)$ be a $2r$-dimensional quadratic space over a field k and let $\mathbf{h}$ be the two-dimensional isotropic quadratic space. Then $\mathrm{G} = \mathrm{Spin}_f$ is quasi-split over k if and only if $(\mathbf{V}, f) \simeq \mathbf{h}^{r-1} \oplus D$ for a two-dimensional space D, and G is split or not according to $D \simeq \mathbf{h}$ or not (see [4, §23.4] where the result is explained for SO_f).

We remind that when k is a number field, quadratic forms of a given degree are classified by looking at every local extension k_v of k (*the Hasse-Minkowski theorem*). Quadratic forms over non-archimedean extensions are classified by the discriminant (denoted by δ) and the *Hasse symbol*. We refer to O'Meara's monograph [22] for the facts about quadratic forms. By the Hasse symbol we mean the normalization of the Hasse invariant which is used in this book.

The group $\mathrm{G} = \mathrm{Spin}_f$ over a non-archimedean local field k_v has the following structure:

(1) G is a split inner form if and only if $\delta(f)$ is equal to $\delta(\mathbf{h}^r) = (-1)^r$ up to an element of $(k_v^*)^2$. In this case G is split exactly when f has the same Hasse symbol as $\mathbf{h}^r$.

(2) For other values of $\delta(f)$, G is quasi-split but non-split and $\delta(f) \in k_v^*/(k_v^*)^2$ determines uniquely the k_v-isomorphism class of G.

3.3. Recall that in odd hyperbolic dimension, besides the construction based on quadratic forms, there exist (cocompact) arithmetic lattices defined by skew-hermitian forms over division quaternion algebras (see [16, Sec. 2] for the details of this construction). Moreover, Cayley algebras produce examples of arithmetic subgroups related to triality forms ${}^{3,6}\mathrm{D}_4$. If we exclude this latter case (which concerns only the hyperbolic dimension seven), we can use the language of algebras with involution to unify the description of admissible groups (we refer to [11] for the notions related to this subject). Namely, all non-trialitarian k-groups which are admissible in the sense of Section 2.2 have (up to k-isomorphism) the form $\mathrm{Spin}(A,\sigma)$, where A is a central simple algebra and σ is an orthogonal involution. More precisely, A is of the form $\mathrm{End}_R(M)$ for a quaternion algebra R over k and an R-module M. If the admissible group is defined by a skew-hermitian form h, then R corresponds to the division algebra over which h is defined [11, Theorem 4.2]. On the other hand, the algebra R *splits*, i.e. $R \simeq \mathrm{End}_k(k^2)$, exactly when A has the form $\mathrm{End}_k(\mathbf{V})$ for a quadratic space $(\mathbf{V}, f)$ with σ being the associated adjoint involution. In this case $\mathrm{Spin}(A,\sigma)$ naturally identifies with the k-group Spin_f. The following result can be considered as a variation on the Hasse-Minkowski theorem.

Lemma 3.4. *Let* G *be an admissible k-group of type different from* ${}^{3,6}\mathrm{D}_4$. *Suppose that for each place $v \in V_\infty \cup V_f$,* G *is k_v-isomorphic to some group* Spin *of a quadratic form over k_v. Then* G *is k-isomorphic to* Spin_f *for some quadratic form f over k.*

Proof. By Section 3.3, the group G has the form $\mathrm{Spin}(\mathrm{End}_R(M),\sigma)$ for some quaternion algebra R over k. By the hypothesis R splits over each completion k_v of the field k. This implies that R splits already over k [17, Theorem 2.7.2]. So G has the form Spin_f for some quadratic form f. □

3.5. Let us now construct an admissible group G_0 which will give rise to a cocompact arithmetic subgroup of small covolume. In this case we have to consider real number fields k different from $\mathbb{Q}$. To get small covolume, by Section 2.6, we need to consider fields k and ℓ whose discriminants have small absolute values. The smallest possible value for $\mathscr{D}_k$ is attained for $k = k_0 = \mathbb{Q}[\sqrt{5}]$. The smallest discriminant of a field ℓ which satisfies the conditions of Proposition 2.5 is $\mathscr{D}_\ell = 275$. It corresponds to a quartic field with a defining polynomial $x^4 - x^3 + 2x - 1$. This pair (k_0, ℓ_0) comes out, for example, with the following quadratic form:

$$f_0(x_0, \ldots, x_n) = \left((-1)^r \cdot 3 - 2\sqrt{5}\right) x_0^2 \,+\, x_1^2 \,+\, \cdots \,+\, x_n^2.$$

Proposition 3.6. *The group* $\mathrm{G}_0 = \mathrm{Spin}_{f_0}$ *is quasi-split at every finite place* $v \in V_f$. *It is the unique admissible group attached to* (k_0, ℓ_0) *which respects this property.*

Proof. The ideal of $\mathscr{O}_{k_0}$ generated by $(-1)^r\delta(f_0)$ is prime. It corresponds to the unique finite place v_r of k_0 at which the extension $\ell_0|k_0$ ramifies. At this ramified place G_0 cannot be inner, and from Section 3.2 it follows that it has to be quasi-split. Let v_2 be the finite place of k_0 corresponding to the prime ideal generated by 2. Then for every finite place v different from v_2 and v_r the fact that the discriminant $\delta(f_0)$ is a unit at v implies that the Hasse invariant is trivial and thus G_0 is quasi-split over k_v. For the remaining place v_2 one can check that the equations $x^2 = \pm\delta(f_0)$ have no solution modulo (8) in the ring of integers of k_0, proving that $\pm\delta(f_0)$ cannot be squares in the local field $(k_0)_{v_2}$. Hence by Section 3.2, G_0 is quasi-split over $(k_0)_{v_2}$ as well.

To prove the uniqueness result we first note that a quasi-split group of type ${}^1\mathrm{D}_r$ or ${}^2\mathrm{D}_r$ is necessarily given by a quadratic form (see [11, Example 27.10]). Since admissibility implies that the group is also defined by quadratic forms over the archimedean completions (namely, by the forms of signature $(n,1)$ and $(n+1,0)$), we can apply Lemma 3.4 to deduce that an admissible group respecting the property stated for the group G_0 in the proposition must be of the type Spin_f, for some quadratic form f defined over k_0. Specifying that Spin_f has the same splitting field ℓ_0 as the group G_0 implies that $\mathrm{Spin}_f \simeq \mathrm{G}_0$ over each completion of k_0. By the Hasse-Minkowski theorem we conclude that such a group Spin_f is k_0-isomorphic to G_0. □

3.7. In order to construct a principal arithmetic subgroup of small covolume we need to consider parahoric subgroups of the maximal volume. When G is quasi-split over k_v and splits over its maximal unramified extension there exist hyperspecial parahoric subgroups in $\mathrm{G}(k_v)$ and those are of the maximal volume. If G does not respect these hypothesis, the parahoric subgroups of maximal volume are the special ones (see [31, Sec. 3.8.2]).

Proposition 3.6 implies the existence of a coherent collection $\mathscr{P}$ of parahoric subgroups $\mathrm{P}_v \subset \mathrm{G}_0(k_v)$ which are hyperspecial for every place v of $k = k_0$ unramified in ℓ_0 and special (but not hyperspecial) for the unique ramified place v_r corresponding to the ideal $(3 - (-1)^r 2\sqrt{5})$. Let us denote by Λ_0 the principal arithmetic subgroup attached to $\mathscr{P}$. Then by Sections 2.6 and 2.7 its covolume is given by:

$$\mu(H/\Lambda_0) = 5^{r^2 - r/2} 11^{r-1/2}\ C(r)^2\ \underbrace{\zeta_k(2)\zeta_k(4)\cdots\zeta_k(2r-2)L_{\ell|k}(r)}_{(*)}. \tag{6}$$

Similarly to the way it was done in [2, proof of Thm. 4.1, p. 760], we can show that the product $(*)$ is bounded by 2. Hence we get

$$\mu(H/\Lambda_0) \quad < \quad 2 \cdot 5^{r^2-r/2}\ 11^{r-1/2}\ C(r)^2\ . \tag{7}$$

For small r we will use a better estimate for the product $(*)$: We can check using Pari/GP (which can compute such an expression up to a given precision) that this product is bounded by 1.17 when $r \leq 16$. Thus for these r we have a better bound:

$$\mu(H/\Lambda_0) \quad < \quad 1.17 \cdot 5^{r^2-r/2}\ 11^{r-1/2}\ C(r)^2\ . \tag{8}$$

These bounds will be used in the proof of Theorem 1 in Sections 7 and 8.

3.8. Now let us consider the non-compact case. It corresponds to $k = \mathbb{Q}$. In contrary to the preceding discussion here we have to distinguish some cases according to the dimension $n = 2r-1$.

First consider r even. By Section 2.3, the field ℓ is a non-trivial extension of $\mathbb{Q}$, and by Proposition 2.5 we see that ℓ must be a quadratic imaginary field. Among these fields the smallest discriminant is attained for $\ell_1 = \mathbb{Q}[\sqrt{-3}]$. Let us take

$$f_1(x_0,\dots,x_n) = -3x_0^2 + x_1^2 + \cdots + x_n^2.$$

For odd r spinor groups of inner type come out. Here we can take

$$f_1(x_0,\dots,x_n) = -x_0^2 + x_1^2 + \cdots + x_n^2.$$

With the quadratic form f_1 defined this way (depending on r) and $\mathrm{G}_1 = \mathrm{Spin}_{f_1}$, the "non-compact" analogue of Proposition 3.6 is given by the following proposition.

Proposition 3.9.

(1) *If r is even, G_1 is quasi-split at every finite place. It is the unique admissible $\mathbb{Q}$-group with the splitting field $\mathbb{Q}[\sqrt{-3}]$ which has this property.*

(2) *For $r = 2m+1$ odd, G_1 is the unique admissible $\mathbb{Q}$-group of type ${}^1\mathrm{D}_r$ which respects the conditions:*
   - (a) *if m is even, G_1 is split over $\mathbb{Q}_p$ for every prime p;*
   - (b) *if m is odd, G_1 is split over $\mathbb{Q}_p$ for every prime $p \neq 2$ and $\mathrm{G}_1|\mathbb{Q}_2$ is a split inner form but non-split.*

Proof. Since the admissible $\mathbb{Q}$-groups are all of the form Spin_f, in all the cases the uniqueness statement follows from the Hasse-Minkowski theorem in a similar way as in the proof of Proposition 3.6. It remains to prove the specific properties of G_1.

When r is even the situation is entirely similar to that in Proposition 3.6: the Hasse invariant ensures that G_1 is quasi-split at primes $p \neq 2,3$, and as $\ell_1|\mathbb{Q}$ is ramified at 3, G_1 is also quasi-split over $\mathbb{Q}_3$. The equation $x^2 = -3$ has no solutions in $\mathbb{Q}_2$ therefore G_1 is quasi-split over $\mathbb{Q}_2$ as well.

For r odd, G_1 is inner at every place of $\mathbb{Q}$. The Hasse invariant is trivial at each prime $p \neq 2$. Computing the Hasse invariant of f_1 and $\mathbf{h}^r$ over $\mathbb{Q}_2$ leads to the different results for m even or odd, which give the second part of the proposition. □

This proposition implies the existence of a coherent collection $\mathscr{P}$ of parahoric subgroups P_v which are all hyperspecial except for at most one place $v = (p)$ of $\mathbb{Q}$, where we choose P_v to be special. For r even this exception is attached to the prime $p = 3$ which is ramified in ℓ_1. For $r = 2m+1$: if m is odd, P_v is not hyperspecial for $v = (2)$, whereas for m even all parahoric subgroups can be chosen to be hyperspecial.

Let Λ_1 be the principal arithmetic subgroup of H determined by $\mathscr{P}$. In Table 1 we list the covolumes of Λ_1 for the different cases. Note that in the last case the value of $\lambda_{(2)}$ (see Section 2.7) is non-trivial. It can be computed using the description of the reductive group $\overline{\mathrm{M}}_{(2)}$ attached to the special parahoric subgroup $\mathrm{P}_{(2)}$. This description is obtained from the local index of $\mathrm{G}(\mathbb{Q}_2)$, which according to Tits [31, Sec. 4.3] is ${}^2\mathrm{D}_r'$ (resp. ${}^2\mathrm{A}_3$ for $r = 3$). More generally, we can compute that if $\mathrm{G}(\mathbb{Q}_p)$ is of this local type, then the lambda factor coming from a special parahoric subgroup is given by:

r	splitting field ℓ	$\mu(H/\Lambda_1)$
r even	$\mathbb{Q}[\sqrt{-3}]$	$3^{r-1/2}\, C(r)\zeta(2)\zeta(4)\cdots\zeta(2r-2)L_{\ell\vert\mathbb{Q}}(r)$
$r \equiv 1 \bmod 4$	$\mathbb{Q}$	$C(r)\zeta(2)\zeta(4)\cdots\zeta(2r-2)\zeta(r)$
$r \equiv 3 \bmod 4$	$\mathbb{Q}$	$\lambda_{(2)}C(r)\zeta(2)\zeta(4)\cdots\zeta(2r-2)\zeta(r)$

TABLE 1. Covolume of Λ_1

$$\lambda_{(p)} \quad = \quad \frac{(p^r-1)\left(p^{r-1}-1\right)}{p+1}. \tag{9}$$

3.10. Using Prasad's volume formula it can be shown without too much effort that Λ_0 (resp. Λ_1) is of minimal covolume in the class of *principal* cocompact (resp. non-cocompact) arithmetic subgroups of H. However, these groups are not necessarily maximal arithmetic subgroups of H. Their normalizers

$$\Gamma_0 := N_H(\Lambda_0) \quad \text{and} \quad \Gamma_1 := N_H(\Lambda_1)$$

are maximal (see Section 2.8) and they will turn out to be the hyperbolic arithmetic lattices of minimal covolume. In the next section we will explain how the computation of the index $[N_H(\Lambda) : \Lambda]$ (with a principal arithmetic subgroup Λ) can be carried out. General estimates obtained there are used in the proofs of Theorems 1 and 2. In Section 5 we deal specifically with the computation of $[\Gamma_0 : \Lambda_0]$ and $[\Gamma_1 : \Lambda_1]$.

4. ESTIMATING THE INDEX $[\Gamma : \Lambda]$

We use the notation of Section 2. In particular, G is an admissible group defined over a totally real number field k and Λ denotes a principal arithmetic subgroup of $H = \mathrm{G}(k_{v_0})$ with a normalizer Γ.

4.1. Let C be the center of G and $\overline{\mathrm{G}}$ be the adjoint group. We have an exact sequence of k-isogenies:

$$1 \to \mathrm{C} \to \mathrm{G} \xrightarrow{\phi} \overline{\mathrm{G}} \to 1.$$

This induces an exact sequence in Galois cohomology (see [24, Sec. 2.2.3]):

$$\mathrm{G}(K) \xrightarrow{\phi} \overline{\mathrm{G}}(K) \xrightarrow{\delta} \mathrm{H}^1(K,\mathrm{C}) \to \mathrm{H}^1(K,\mathrm{G}), \tag{10}$$

valid for every field extension $K|k$.

The group G is simply connected hence for each $v \in V_f$ we have $\mathrm{H}^1(k_v,\mathrm{G}) = 1$ (see [24, Theorem 6.4]). The group $\overline{\mathrm{G}}(k_v)$ acts by conjugation on the set of parahoric subgroups and the action of $\phi(\mathrm{G}(k_v))$ is trivial, hence this induces a homomorphism

$$\xi_v : \mathrm{H}^1(k_v,\mathrm{C}) \to \mathrm{Aut}(\Delta_v).$$

The image of ξ_v is denoted by Ξ_v, and we can define the map

$$\xi : \mathrm{H}^1(k,\mathrm{C}) \to \bigoplus_{v\in V_f} \Xi_v$$

as a product of all ξ_v (restricted to $\mathrm{H}^1(k,\mathrm{C})$). This material is explained in more detail in [5, Sec. 2].

4.2. Given $E \le \mathrm{H}^1(k,\mathrm{C})$, we denote by E_ξ its subgroup which acts trivially on each Δ_v and by E_θ the subgroup which stabilizes the type $\theta = \prod_{v\in V_f} \theta_v$ attached to Λ as in Section 2.2. We have

$$\#E_\theta \le \#E_\xi \prod_{v\in V_f} \#\Xi_{\theta_v}, \tag{11}$$

where Ξ_{θ_v} is the subgroup of Ξ_v stabilizing θ_v.

Now by [26], [5, Prop. 2.9], we have an exact sequence

$$1 \to \mathrm{C}(k_{v_0})/\left(\mathrm{C}(k)\cap\Lambda\right) \to \Gamma/\Lambda \to \delta(\overline{\mathrm{G}}(k))'_\theta \to 1, \tag{12}$$

where

$$\begin{aligned} \delta(\overline{\mathrm{G}}(k))' &= \delta(\overline{\mathrm{G}}(k)) \cap (\delta\circ\phi)(\mathrm{G}(k_{v_0})), \text{ and} \\ \delta(\overline{\mathrm{G}}(k))'_\theta &= \delta(\overline{\mathrm{G}}(k))' \cap \mathrm{H}^1(k,\mathrm{C})_\theta. \end{aligned}$$

In our case Λ, being a principal arithmetic subgroup of $\mathrm{G}(k)$, contains the center $\mathrm{C}(k)$. Also, $\mathrm{C}(k_{v_0}) = \{\pm 1\}$ and using the description of C in Section 4.4 we can check that if G is not of type ${}^{3,6}\mathrm{D}_4$, then $\mathrm{C}(k) = \mathrm{C}(k_{v_0})$. This implies that for these types $\Gamma/\Lambda \simeq \delta(\overline{\mathrm{G}}(k))'_\theta$. For G of type ${}^{3,6}\mathrm{D}_4$ we have $\mathrm{C}(k) = 1$ and the quotient Γ/Λ is a $\mathbb{Z}/2\mathbb{Z}$-extension of $\delta(\overline{\mathrm{G}}(k))'_\theta$.

4.3. In [5, Sec. 3] a principal arithmetic subgroup $\Lambda^{\mathfrak{m}}$ commensurable with Λ is chosen in such a way that the *generalized index* of Λ in $\Lambda^{\mathfrak{m}}$ respects the following inequality:

$$[\Lambda^{\mathfrak{m}} : \Lambda] = \frac{[\Lambda^{\mathfrak{m}} : \Lambda^{\mathfrak{m}} \cap \Lambda]}{[\Lambda : \Lambda^{\mathfrak{m}} \cap \Lambda]} \geq \prod_{v \in V_f} \#\Xi_{\theta_v}. \tag{13}$$

We let $\theta^{\mathfrak{m}} = (\theta_v^{\mathfrak{m}})$ be the global type of the principal subgroup $\Lambda^{\mathfrak{m}}$. By the construction in [5], the types $\theta_v^{\mathfrak{m}}$ are all special which implies that for every $v \in V_f$ we have $\Xi_{\theta_v^{\mathfrak{m}}} = 1$. It follows that $\delta(\overline{\mathrm{G}}(k))'_{\theta^{\mathfrak{m}}} = \delta(\overline{\mathrm{G}}(k))'_{\xi}$. As this value depends only on the group G, we conclude that the covolume of the normalizer $\Gamma^{\mathfrak{m}} = N_H(\Lambda^{\mathfrak{m}})$ depends only on G.

Now we have

$$\mu(H/\Gamma) = \frac{\mu(H/\Lambda)}{[\Gamma : \Lambda]} = \frac{\mu(H/\Lambda^{\mathfrak{m}})[\Lambda^{\mathfrak{m}} : \Lambda]}{[\Gamma : \Lambda]}.$$

By (13), it gives

$$\mu(H/\Gamma) \geq \frac{\mu(H/\Lambda^{\mathfrak{m}})}{[\Gamma : \Lambda]} \prod_{v \in V_f} \#\Xi_{\theta_v}. \tag{14}$$

From the other hand, by (11) and (12),

$$[\Gamma : \Lambda] \leq [\Gamma^{\mathfrak{m}} : \Lambda^{\mathfrak{m}}] \prod_{v \in V_f} \#\Xi_{\theta_v}. \tag{15}$$

Combining inequalities (14) and (15) we obtain

$$\mu(H/\Gamma) \geq \frac{1}{[\Gamma^{\mathfrak{m}} : \Lambda^{\mathfrak{m}}]} \mu(H/\Lambda^{\mathfrak{m}}) = \mu(H/\Gamma^{\mathfrak{m}}).$$

This means that $\Gamma^{\mathfrak{m}}$ is of the smallest covolume among the arithmetic subgroups attached to G. Its covolume depends on $\#\delta(\overline{\mathrm{G}}(k))'_{\xi}$ and we will now focus on the computation of this order.

In the rest of the article we will write $\Lambda = \Lambda^{\mathfrak{m}}$ to indicate that Λ is commensurable with $\Lambda^{\mathfrak{m}}$ and has a similar local structure, i.e., $\theta^{\mathfrak{m}}$ is in the $\overline{\mathrm{G}}(\mathbb{A}_f)$-orbit of θ (cf. Section 6.1). Similarly, the notation $\Gamma^{\mathfrak{m}} = \Gamma$ will be used for their normalizers. The equality is an abuse of notation. However, it is clear that in this case Γ (resp. Λ) has the same covolume as $\Gamma^{\mathfrak{m}}$ (resp. $\Lambda^{\mathfrak{m}}$).

4.4. We remind here the description of the center C from [24, Table on p. 332]. Let $\mathfrak{n} = 4$, $\varepsilon = 1$ if G is of absolute type D_{2m+1}, and $\mathfrak{n} = 2$, $\varepsilon = 2$, otherwise. The inner form of type ${}^1\mathrm{D}_r$ has as its center the group $\mu_{\mathfrak{n}}^{\varepsilon}$ of order $\mathfrak{n}^{\varepsilon}$. For the type ${}^2\mathrm{D}_{2m}$ the center C is isomorphic to $\mathrm{R}_{\ell|k}(\mu_2)$, where $\mathrm{R}_{\ell|k}$ denotes the *restriction of scalars* relative to the field extension $\ell|k$ defined in Section 2.4. In the remaining cases (${}^2\mathrm{D}_{2m+1}$ and ${}^{3,6}\mathrm{D}_4$) the center is isomorphic to the kernel $\mathrm{R}^{(1)}_{\ell|k}(\mu_{\mathfrak{n}})$ of the *norm map* $N_{\ell|k} : \mathrm{R}_{\ell|k}(\mu_{\mathfrak{n}}) \to \mu_{\mathfrak{n}}$. Here we have the following exact sequence:

$$1 \to \mathrm{R}^{(1)}_{\ell|k}(\mu_{\mathfrak{n}}) \to \mathrm{R}_{\ell|k}(\mu_{\mathfrak{n}}) \overset{N_{\ell|k}}{\to} \mu_{\mathfrak{n}} \to 1. \tag{16}$$

4.5. From Section 4.4 we can deduce a description of $\mathrm{H}^1(k,\mathrm{C})$ for the admissible groups. For the type ${}^1\mathrm{D}_{2m+1}$, we have $\mathrm{C} = \mu_4$ and $\mathrm{H}^1(k,\mathrm{C}) = k^*/(k^*)^4$. If G is of type ${}^2\mathrm{D}_{2m}$, then $\mathrm{H}^1(k,\mathrm{C}) = \mathrm{H}^1(k,\mathrm{R}_{\ell|k}(\mu_2)) = \ell^*/(\ell^*)^2$. Dealing with the remaining cases requires considering the cohomological exact sequence associated to (16), which gives us the following exact sequence:

$$1 \to \mu_{\mathfrak{n}}(k)/N_{\ell|k}(\mu_{\mathfrak{n}}(\ell)) \to \mathrm{H}^1(k,\mathrm{C}) \to \ker\left(\ell^*/(\ell^*)^{\mathfrak{n}} \overset{N_{\ell|k}}{\to} k^*/(k^*)^{\mathfrak{n}}\right) \to 1. \tag{17}$$

By Proposition 2.5(iii), the $\mathfrak{n}$-roots of unity in k and ℓ are ± 1 with the only possible exception when $k = \mathbb{Q}$ and $\ell = \mathbb{Q}[\sqrt{-1}]$. Therefore, the second group in (17) is $\{\pm 1\}$ if G is of type ${}^2\mathrm{D}_{2m+1}$ and is trivial for ${}^{3,6}\mathrm{D}_4$.

In all the cases we can define the *image* of $\mathrm{H}^1(k,\mathrm{C})$ in the group $\ell^*/(\ell^*)^{\mathfrak{n}}$ (for the inner types we have $\ell = k$). To describe this image we set $\mathbf{L} = \ell^*$ in the cases ${}^2\mathrm{D}_{2m}$ and ${}^1\mathrm{D}_{2m+1}$, and $\mathbf{L} = \left\{x \in \ell^* \mid N_{\ell|k}(x) \in (k^*)^{\mathfrak{n}}\right\}$ in the remaining cases ${}^2\mathrm{D}_{2m+1}$ and ${}^{3,6}\mathrm{D}_4$. Then, except for the case ${}^2\mathrm{D}_{2m+1}$, $\mathbf{L}/(\ell^*)^{\mathfrak{n}}$ is an isomorphic image of $\mathrm{H}^1(k,\mathrm{C})$. In the case ${}^2\mathrm{D}_{2m+1}$, $\mathrm{H}^1(k,\mathrm{C})$ is a $\mathbb{Z}/2\mathbb{Z}$-extension of $\mathbf{L}/(\ell^*)^{\mathfrak{n}}$.

4.6. Let $A = \delta(\overline{\mathrm{G}}(k))'$ and let $\mathbf{A} \leq \mathbf{L}$ be such that the image of A in $\mathbf{L}/(\ell^*)^{\mathfrak{n}}$ is equal to $\mathbf{A}/(\ell^*)^{\mathfrak{n}}$. In order to describe the group A we need to solve two problems: the first is related to understanding the image of $\overline{\mathrm{G}}(k)$ and the second is considering the image of $\mathrm{G}(k_{v_0})$ (see Section 4.2). Both of these images can be described by looking at the archimedean places.

To begin with, let us consider the whole image of $\delta : \overline{\mathrm{G}}(k) \to \mathrm{H}^1(k,\mathrm{C})$. By (10) it is equal to $\ker(\mathrm{H}^1(k,\mathrm{C}) \to \mathrm{H}^1(k,\mathrm{G}))$. The Hasse principle for G implies that $\mathrm{H}^1(k,\mathrm{G})$ is isomorphic to $\prod_{v \in V_\infty} \mathrm{H}^1(k_v,\mathrm{G})$ [24, Theorem 6.6]. Therefore,

$$\delta(\overline{\mathrm{G}}(k)) = \ker\left(\mathrm{H}^1(k,\mathrm{C}) \to \textstyle\prod_{v \in V_\infty} \mathrm{H}^1(k_v,\mathrm{G})\right). \tag{18}$$

Now let us restrict our attention to the group A, which means that we have to add the condition that the elements are in the image of $\mathrm{G}(k_{v_0})$ under the map $\delta \circ \phi$. Using (10) with $K = k_{v_0}$, we see that these elements are exactly those which are trivial in $\mathrm{H}^1(k_{v_0},\mathrm{C})$. For the other infinite places $v \neq v_0$ the group $\mathrm{G}(k_v) \simeq \mathrm{Spin}(n+1)$ maps surjectively onto $\overline{\mathrm{G}}(k_v)$ and therefore $\ker(\mathrm{H}^1(k_v,\mathrm{C}) \to \mathrm{H}^1(k_v,\mathrm{G})) = 1$. Thus from (18) we get:

$$A = \ker\left(\mathrm{H}^1(k,\mathrm{C}) \to \textstyle\prod_{v \in V_\infty} \mathrm{H}^1(k_v,\mathrm{C})\right). \tag{19}$$

Let us write $V_\infty = \{v_0, v_1, \ldots\}$ for the infinite places of k. Suppose that G is not of type ${}^{3,6}\mathrm{D}_4$. Then the tensor product $\ell_{v_i} = \ell \otimes k_{v_i}$ is isomorphic to $\mathbb{R}$ if $\ell = k$ and isomorphic either to $\mathbb{C}$ or $\mathbb{R} \oplus \mathbb{R}$ according to the type of G over k_{v_i} when $\ell|k$ is a quadratic extension. In the latter case, let us write σ_i and σ_i' for the two real embeddings of ℓ extending the place v_i. In particular, for $x \in \ell$, the notation $x^{\sigma_i} > 0$ or $x^{\sigma_i'} > 0$ will make sense.

Proposition 4.7. *We have the following description of A:*

- *(1) For $k \neq \mathbb{Q}$ we have $A \simeq \mathbf{A}/(\ell^*)^{\mathfrak{n}}$, and, in particular,*
  - *(a) if G is of type ${}^2\mathrm{D}_{2m+1}$, $\mathbf{A} = \{x \in \mathbf{L} \mid x^{\sigma_0} > 0\}$;*
  - *(b) if G is of type ${}^2\mathrm{D}_{2m}$, $\mathbf{A} = \left\{x \in \ell^* \mid x^{\sigma_i}, x^{\sigma_i'} > 0 \ \forall i \neq 0\right\}$.*
- *(2) For $k = \mathbb{Q}$:*
  - *(a) if G is of type ${}^1\mathrm{D}_{2m+1}$, $\mathbf{A} = \{x \in \mathbb{Q}^* \mid x > 0\}$ and $A \simeq \mathbf{A}/(\mathbb{Q}^*)^4$;*
  - *(b) if G is of type ${}^2\mathrm{D}_{2m+1}$, A is a $\mathbb{Z}/2\mathbb{Z}$-extension of $\mathbf{A}/(\ell^*)^4$, where $\mathbf{A} = \{x \in \mathbf{L} \mid x^{\sigma_0} > 0\}$;*
  - *(c) if G is of type ${}^2\mathrm{D}_{2m}$, $\mathbf{A} = \ell^*$ and $A \simeq \mathbf{A}/(\ell^*)^2$.*

Proof. By (19), an element of $\mathrm{H}^1(k, \mathrm{C})$ is in A if and only if its image in $\mathrm{H}^1(k_v, \mathrm{C})$ is trivial for every $v \in V_\infty$. The group $\mathrm{H}^1(k, \mathrm{C})$ is described in terms of the field ℓ and the mapping $\mathrm{H}^1(k, \mathrm{C}) \to \mathrm{H}^1(k_v, \mathrm{C})$ can be understood when considering the inclusion $\ell \to \ell_v$. Let us begin with the type ${}^2\mathrm{D}_{2m+1}$. In this case $\ell_{v_0} \simeq \mathbb{R} \oplus \mathbb{R}$ and $\ell_{v_i} \simeq \mathbb{C}$ for $i \neq 0$. The exact sequence (17) takes the following form over v_0, resp. v_i:

$$\begin{array}{lccccccccc} (20) & 1 & \to & 1 & \to & \mathrm{H}^1(k_{v_0}, \mathrm{C}) & \to & \left\{(\mathbb{R}_{>0})^2, (\mathbb{R}_{<0})^2\right\} & \to & 1; \\ (21) & 1 & \to & \{\pm 1\} & \to & \mathrm{H}^1(k_{v_i}, \mathrm{C}) & \to & 1 & \to & 1. \end{array}$$

Here $(\mathbb{R}_{>0})^2$ and $(\mathbb{R}_{<0})^2$ are meant as the elements of $\ell_{v_0}^*/(\ell_{v_0}^*)^4 \simeq (\mathbb{R}^*/\mathbb{R}_{>0})^2$. If $k \neq \mathbb{Q}$, there exists a place $v_i \neq v_0$ and hence by (21) the $\mathbb{Z}/2\mathbb{Z}$-extension is killed. By (20), an element $x \in \mathbf{A}$ must respect: $x^{\sigma_0}, x^{\sigma_0'} > 0$. But since the norm of x must be positive (to be in $\mathbf{L}$), $x^{\sigma_0} > 0$ implies positivity of $x^{\sigma_0'}$. These facts together give us *(1a)* and *(2b)*.

For the type ${}^1\mathrm{D}_{2m+1}$ (arising only when $k = \mathbb{Q}$, so $V_\infty = \{v_0\}$), we have $\mathrm{H}^1(k_{v_0}, \mathrm{C}) = \mathrm{H}^1(\mathbb{R}, \mu_4) \simeq \mathbb{R}^*/(\mathbb{R}^*)^2$. An element $x \in \mathbb{Q}^*$ has then trivial image in $\mathrm{H}^1(k_{v_0}, \mathrm{C})$ exactly when x is positive. This proves *(2a)*. In the case ${}^2\mathrm{D}_{2m}$, the group G becomes inner over k_{v_0}, which means that $\ell_{v_0} \simeq \mathbb{C}$. Then $\mathrm{H}^1(k_{v_0}, \mathrm{C})$ is trivial and the condition at v_0 is empty. Hence for $k = \mathbb{Q}$ we get *(2c)*. For $k \neq \mathbb{Q}$ we need to include the conditions at v_i for $i \neq 0$. As $\mathrm{H}^1(k_{v_i}, \mathrm{C}) \simeq (\mathbb{R}^*/\mathbb{R}_{>0})^2$, this gives *(1b)*. □

4.8. We need to fix some more notations. Let T_1 denote the set of places $v \in V_f$ such that if G is of inner type over k_v, it does not split, and if it is of outer type then it is not quasi-split over k_v but splits over its maximal unramified extension $\hat{k}_v$. In the latter case let R be the subset of $V_f \setminus T_1$ such that G does not split over $\hat{k}_v$, which can be also described as the set of places of k which are ramified in ℓ/k. We have $T_1 \subset T$, where T is defined as in Section 2.7 for some principal arithmetic subgroup Λ of G.

Let S be a subset of V_f. For the subgroup $\mathbf{A}$ of ℓ^* we denote by $\mathbf{A}_{\mathfrak{n}}^S$ its subgroup which consists of the elements x such that $\tilde{v}(x) \in \mathfrak{n}\mathbb{Z}$ for every *normalized* nonarchimedean valuation $\tilde{v}$ of ℓ which is not above some place from S. If $S = \emptyset$ we simply write $\mathbf{A}_{\mathfrak{n}}$. We introduce a refinement of the notation from Section 4.2: for the finite subset $S \subset V_f$ we denote by $A_{\xi,S}$ the subgroup of A which acts trivially on Δ_v for every $v \notin S$. Clearly, $A_{\xi,\emptyset} = A_\xi$.

4.9. By Sections 4.2–4.3 the index $[\Gamma:\Lambda]$ depends on the order of A_ξ. We can bound this order using results of [5]. For $v \notin S = R \cup T_1$, an element of A acts trivially on Δ_v if and only if its image x in $\mathbf{A}$ has $\tilde{v}(x) \in \mathfrak{n}\mathbb{Z}$ for every normalized valuation $\tilde{v}$ of ℓ extending v [5, Lemma 2.3 and Prop. 2.7]. This implies that the image of $A_{\xi,S}$ in $\mathbf{A}/(\ell^*)^{\mathfrak{n}}$ is given by $\mathbf{A}_{\mathfrak{n}}^S/(\ell^*)^{\mathfrak{n}}$. Therefore, we can describe the relation between A_ξ and $\mathbf{A}_{\mathfrak{n}}/(\ell^*)^{\mathfrak{n}}$ by the following diagram:

$$\begin{tikzcd}
A_{\xi,S} \arrow[r, "\bar{q}"] & \mathbf{A}_{\mathfrak{n}}^S/(\ell^*)^{\mathfrak{n}} \\
A_\xi \arrow[u, hook, "q'"] & \mathbf{A}_{\mathfrak{n}}/(\ell^*)^{\mathfrak{n}} \arrow[u, hook, "q"]
\end{tikzcd}$$

The integers q and q' stand for the indices of the vertical inclusions. The order of the kernel of the horizontal map is denoted by $\bar{q}$. By Proposition 4.7, $\bar{q}$ is equal to 1 with the only exception of the type ${}^2\mathrm{D}_{2m+1}$ with $k=\mathbb{Q}$, where $\bar{q}=2$. It is possible to compute the order of $\mathbf{A}_{\mathfrak{n}}/(\ell^*)$ using the approach which we will explain in Section 4.11. The indices q and q' are difficult to evaluate precisely (in particular, for this we would need a counterpart of [5, Prop. 2.7] for $v \in S$). However, it is possible to bound their values:

Proposition 4.10. *The indices q and q' satisfy*

(1) (a) $q \leq 4^{\#T_1}$, if G is of absolute type D_{2m+1};
(b) $q \leq 2^{\#R}4^{\#T_1}$, if G is of type ${}^2\mathrm{D}_{2m}$ or ${}^{3,6}\mathrm{D}_4$.
(2) q' is a factor of $\prod_{v\in S} \#\Xi_v$.

Proof. The proof of the first assertion uses the same arguments as in [5, Sec. 5], where the authors consider the whole $\mathrm{H}^1(k,\mathrm{C})$ instead of our group A. More precisely, for the inner type ${}^1\mathrm{D}_{2m+1}$ we obtain the bound using the argument of [5, Sec. 5.1]. For the other cases we use the same idea as in the proof of Lemma 5.4 from [5]. The second assertion is obvious since $A_{\xi,S}$ is mapped into $\prod_{v\in S}\Xi_v$ with the kernel A_ξ. □

4.11. We denote by U_ℓ the group of units of the ring of integers of ℓ and let $U_{\mathbf{A}} = \mathbf{A} \cap U_\ell$. The symbol $\mathscr{P}_\ell$ stands for the principal fractional ideal group of ℓ, $\mathscr{C}_\ell$ denotes the class group and its order (the *class number*) is denoted by h_ℓ. The exact sequence (1) in the proof of [5, Prop. 0.12] can be adapted to our setting by replacing ℓ^* by $\mathbf{A}$. This gives an exact sequence:

$$1 \to U_{\mathbf{A}} \to \mathbf{A} \to \mathscr{P}_{\mathbf{A}} \to 1,$$

where $\mathscr{P}_{\mathbf{A}}$ is a certain subgroup of $\mathscr{P}_\ell$. In all cases we have $(\ell^*)^{\mathfrak{n}} \subset \mathbf{A}$, so that we can take this sequence modulo $(\ell^*)^{\mathfrak{n}}$. Then, following the same line of argument as in the above mentioned proof, we get an exact sequence:

$$1 \to U_{\mathbf{A}}/U_\ell^{\mathfrak{n}} \to \mathbf{A}_{\mathfrak{n}}/(\ell^*)^{\mathfrak{n}} \to \mathscr{C}_{\mathbf{A}} \to 1, \tag{22}$$

where $\mathscr{C}_{\mathbf{A}}$ is a subgroup of $\mathscr{C}_\ell$ (it can be described explicitly, however, we do not need it here).

Proposition 4.12. *The index of $\Lambda = \Lambda^{\mathfrak{m}}$ in its normalizer Γ is bounded by the following value:*

- *(1) For $k \neq \mathbb{Q}$:*
  - *(a) If G is of type ${}^2\mathrm{D}_{2m+1}$, $[\Gamma : \Lambda] \leq 2^{d+1} 4^{\# T_1} h_\ell$;*
  - *(b) If G is of type ${}^2\mathrm{D}_{2m}$, $[\Gamma : \Lambda] \leq 2^{2d-1} 2^{\# R} 4^{\# T_1} h_\ell$;*
  - *(c) If G is of type ${}^{3,6}\mathrm{D}_4$, $[\Gamma : \Lambda] \leq 2^{3d+1} 2^{\# R} 4^{\# T_1} h_\ell$.*
- *(2) For $k = \mathbb{Q}$:*
  - *(a) If G is of type ${}^1\mathrm{D}_{2m+1}$, $[\Gamma : \Lambda] \leq 4^{\# T_1}$;*
  - *(b) If G is of type ${}^2\mathrm{D}_{2m+1}$, $[\Gamma : \Lambda] \leq 8 \cdot 4^{\# T_1} h_\ell$;*
  - *(c) If G is of type ${}^2\mathrm{D}_{2m}$, $[\Gamma : \Lambda] \leq 4 \cdot 2^{\# R} 4^{\# T_1} h_\ell$.*

Proof. Let us deal with the first case. Here we have $[\Gamma : \Lambda] = \# A_\xi$. Proposition 4.10 and the exact sequence (22) allow us to bound the order of A_ξ by $4^{\# T_1} \cdot h_\ell \cdot \#(U_{\mathbf{A}}/U_\ell^4)$. By Dirichlet's units theorem, U_ℓ is a semi-direct product of $\mu(\ell) = \{\pm 1\}$ and $\mathbb{Z}^d$ (where $d+1$ is the number of infinite places of ℓ, as given in Proposition 2.5). Now we use $U_{\mathbf{A}} \subset U_{\mathbf{L}}$ and the fact that the norm map $N_{\ell|k}$ covers U_k^2 (taking x as a preimage of x^2) to get the following bound:

$$\#\left(U_{\mathbf{L}}/U_\ell^4\right) \;\leq\; \frac{\#\left(U_\ell/U_\ell^4\right)}{\#\left(U_k^2/U_k^4\right)} = \frac{2 \cdot 4^d}{2^{d-1}}.$$

Finally, restricting our attention to $U_{\mathbf{A}}$ instead of $U_{\mathbf{L}}$, the description of $\mathbf{A}$ in Proposition 4.7 implies that we have to consider only positive elements in $U_{\mathbf{L}}$, and so we divide the bound by 2. This gives us *(1a)*. The statement *(2b)* is obtained by the same argument with $d = 1$, the only difference comes from the value of q' (2 instead of 1 when $k = \mathbb{Q}$).

The remaining cases use only the bound for q in Proposition 4.10, the bound for $U_\ell/U_\ell^{\mathfrak{n}}$ provided by Dirichlet's units theorem and Proposition 2.5. Note that for triality forms we must keep in mind that the index $[\Gamma : \Lambda]$ is twice the order of A_ξ. □

4.13. Let us point out that in the proof of Prop. 1.3 [2, Add.] it should be written $\mathrm{Im}(\delta)_\theta$ instead of $\mathrm{Im}(\delta)$ everywhere, moreover, in a half of the cases there one has to consider more carefully the set T of places over which the group is non-split. This, however, does not change the result.

5. COMPUTATION OF INDICES $[\Gamma_0 : \Lambda_0]$ AND $[\Gamma_1 : \Lambda_1]$

In this section we compute the indices $[\Gamma_0 : \Lambda_0]$ and $[\Gamma_1 : \Lambda_1]$. The subgroups involved were defined in Section 3. The computation is based on the material of Section 4. Similar kind of arguments have been used in [19], where the authors study groups which act transitively on vertices of Bruhat-Tits buildings.

5.1. Let $k = k_0 = \mathbb{Q}[\sqrt{5}]$ be the field of definition of Λ_0 and let $\ell = \ell_0$ be the splitting field attached to Λ_0. We have $\mathscr{D}_\ell = \mathscr{D}_{\ell_0} = 275$. By construction, the associated group G is of type ${}^2\mathrm{D}_r$ and we have $\Lambda_0 = \Lambda_0^{\mathfrak{m}}$ which implies that $[\Gamma_0 : \Lambda_0] = \# A_\xi$. By Proposition 4.7, the group A is isomorphic to $\mathbf{A}/(\ell^*)^{\mathfrak{n}}$ and has two different descriptions according to the parity of the rank. Furthermore, as in our case $h_\ell = 1$ the exact sequence (22) shows that

$$[\Gamma_0 : \Lambda_0] \quad = \quad \frac{q}{q'}\,\#(U_{\mathbf{A}}/U_\ell^{\mathfrak{n}})\,, \tag{23}$$

where q and q' are the integers defined in Section 4.9.

The two real embeddings of ℓ correspond to the two k-isomorphisms of $k[\sqrt{\alpha}]$ with $\alpha = 3+2\sqrt{5}$. These real embeddings appear over the place v_0 in odd rank (i.e., in this case $\ell_{v_0} = \mathbb{R}\oplus\mathbb{R}$) and over the place v_1 in even rank (i.e., here $\ell_{v_1} = \mathbb{R}\oplus\mathbb{R}$). The field $k[\sqrt{\alpha}]$ has two fundamental units given by

$$\tau_1 = \frac{1+\sqrt{\alpha}}{2}\quad,\quad \tau_2 = \frac{1-\sqrt{\alpha}}{2}.$$

We can identify the group $U_\ell/U_\ell^{\mathfrak{n}}$ with the representative set

$$\left\{\pm\tau_1^i\tau_2^j \mid 0 \leq i,j \leq \mathfrak{n}-1\right\}.$$

The elements which are in $U_{\mathbf{A}}/U_\ell^{\mathfrak{n}}$ are then (cf. Proposition 4.7)

$$\begin{aligned} 1, \tau_1^2\tau_2^2, -\tau_1\tau_2^3, -\tau_1^3\tau_2 \quad &\text{when } r = 2m+1;\\ 1, -\tau_1\tau_2 \quad &\text{when } r = 2m. \end{aligned}$$

Hence:

$$\#(U_{\mathbf{A}}/U_\ell^{\mathfrak{n}}) = \begin{cases} 4 & \text{in case } \mathrm{D}_{2m+1},\\ 2 & \text{in case } \mathrm{D}_{2m}. \end{cases} \tag{24}$$

5.2. We now explain how to determine the values of q and q'. We will make use of the fact stated in [31, Sec. 2.5] that for the ramified place v_{r} (defined in the proof of Proposition 3.6) we have $\#\Xi_{v_{\mathrm{r}}} = 2$, i.e., $\Xi_{v_{\mathrm{r}}}$ is the whole symmetry group of the local Dynkin diagram $\Delta_{v_{\mathrm{r}}}$ of the form



Moreover, we recall that the group G_0 defining Λ_0 is such that $T_1 = \emptyset$.

Let us first assume that $r = 2m$. In this case $\tau_1\sqrt{\alpha} \in \mathbf{A}$ represents an element of $A_{\xi,R}$ which is clearly not in $\mathbf{A}_2/(\ell^*)^2$. This implies that $q = 2$. Moreover, using Hensel's Lemma to detect congruences modulo square in $\ell_{v_{\mathrm{r}}}$, we can check that $-\tau_1\tau_2$ is another element of $A_{\xi,R}$ which is mapped to a non-trivial element of $\mathrm{H}^1(k_{v_{\mathrm{r}}},\mathrm{C}) = \ell_{v_{\mathrm{r}}}^*/(\ell_{v_{\mathrm{r}}}^*)^2$. But since this latter group has order 4 (see [15, Ch. II, Prop. 6]), $A_{\xi,R}$ must map surjectively onto it. In particular, $A_{\xi,R}$ acts non-trivially on $\Delta_{v_{\mathrm{r}}}$ and we get $q' = 2$. By Section 5.1 this implies that $[\Gamma_0 : \Lambda_0] = 2$.

For $r = 2m+1$ by Proposition 4.10 we have $q = 1$. To see that $A_{\xi,R}$ acts effectively on $\Delta_{v_{\mathrm{r}}}$ here we cannot use directly the same argument as for the case $r = 2m$. Indeed, $A_{\xi,R}$ cannot be mapped onto $\mathrm{H}^1(k_{v_{\mathrm{r}}},\mathrm{C})$ since (contrary to the group A) it is an extension by $\mu_4(k_{v_{\mathrm{r}}})/N_{\ell|k}(\mu_4(\ell_{v_{\mathrm{r}}})) = \{\pm 1\}$ of the kernel of the norm map (compare with (17)). But we can check that this subgroup $\{\pm 1\} < \mathrm{H}^1(k_{v_\mathrm{r}}, \mathrm{C})$ acts trivially on Δ_{v_r}. For this we consider the element $(i,i) \in \mu_4(\overline{k}) \times \mu_4(\overline{k}) \simeq \mathrm{R}_{\ell|k}(\mu_4)(\overline{k})$. We have $N_{\ell|k}((i,i)) = i^2 = -1$ and hence from the exact sequence (16) we see (cf. [24, Sec. 1.3.2]) that -1 is mapped to the cocycle $a \in \mathrm{H}^1(k, \mathrm{C})$ given by:

$$\sigma \in \mathrm{Gal}(\overline{k}|k) \quad \mapsto \quad a_\sigma = \begin{cases} -1 & \text{if } \sigma(i) = -i, \\ 1 & \text{otherwise.} \end{cases}$$

We can check that this cocycle class is represented by $\sigma \mapsto {}^\sigma\tilde{g}\tilde{g}^{-1} \in \mathrm{C}$, with $\tilde{g} \in \mathrm{Spin}_{f_0}$ constructed as follows:

$$\tilde{g} \quad = \quad ixy - ixy,$$

for two isotropic vectors $x, y \in (\mathbf{V}, f_0)$ such that $f_0(x+y) = 1/2$. We recall here that Spin_{f_0} is the subgroup of the Clifford group which consists of the elements of norm one which give special orthogonal transformations of $(\mathbf{V}, f_0)$ (see [29, Ch. 9, §3]). A direct computation shows that the image of $\tilde{g}$ in the group SO_{f_0} is given by the matrix $\mathrm{diag}(-1,-1,1,\ldots,1)$, with x, y as the first vectors of the basis completed by orthogonal elements. Therefore, we see that the image $\phi(\tilde{g})$ (see Section 4.1) is in the compact part of any torus of $\overline{\mathrm{G}}(k_{v_\mathrm{r}})$ containing it, which means that the cocycle a acts trivially on Δ_{v_r} (see [31, Sec. 2.5]). This proves that $-1 \in \mu_4(k_{v_\mathrm{r}})$ is trivial in Ξ_{v_r}. Similarly to the case $r = 2m$ we can then check that $A_{\xi,R}$ is mapped onto $\ker\left(\ell_{v_\mathrm{r}}^*/(\ell_{v_\mathrm{r}}^*)^4 \to k_{v_\mathrm{r}}^*/(k_{v_\mathrm{r}}^*)^4\right)$. This shows that $q' = 2$, so that for all ranks we have:

$$[\Gamma_0 : \Lambda_0] \quad = \quad 2. \tag{25}$$

5.3. Now consider the group Γ_1 which is constructed using the group G_1 defined over $\mathbb{Q}$. As in the cocompact case we have $\Lambda_1 = \Lambda_1^\mathrm{m}$. Let first $r = 2m+1$. Then G_1 is of inner type ${}^1\mathrm{D}_{2m+1}$ and by Proposition 4.7 and Section 4.11 we have $\mathbf{A}_4/(\mathbb{Q}^*)^4 = 1$. Now by Proposition 3.9, the set T_1 is empty if m is even, and it follows immediately in this case that $q = q' = 1$. If m is odd, T_1 consists of a single place (2). The elements $2, 4, 8 \in \mathbb{Q}^*$ are three non-trivial elements of $\mathbf{A}$ which determine different elements of A_{ξ,T_1} which are not in $\mathbf{A}_4/(\mathbb{Q}^*)^4$. This allows us to see that $q = 4$. To compute q' we need to identify $2 \in \mathbb{Q}_2^*/(\mathbb{Q}_2^*)^4$ with an element of $\overline{\mathrm{G}}(\mathbb{Q}_2)$. For this consider a basis $x_1, y_1, \ldots, x_r, y_r$ of $(\mathbf{V}|\mathbb{Q}_2, f_1)$ which satisfies the following conditions:

(1) for $j = 1, \ldots, r-2$, x_j and y_j are isotropic vectors with $f_1(x_j + y_j) = 1/2$;
(2) for $j = r-1$ and $j = r$, x_j and y_j are chosen orthonormal.

Such a choice of the basis is possible according to the structure of the space $(\mathbf{V}, f_1)$ over $\mathbb{Q}_2$ (cf. Proposition 3.9). With respect to this basis the matrix

$$\frac{1}{\sqrt{2}}\begin{pmatrix} 2 & 0 & & & & & & & \\ 0 & 1 & & & & & & & \\ & & \ddots & & & & & & \\ & & & 2 & 0 & & & & \\ & & & 0 & 1 & & & & \\ & & & & & 1 & 1 & & \\ & & & & & -1 & 1 & & \\ & & & & & & & 1 & 1 \\ & & & & & & & -1 & 1 \end{pmatrix}$$

gives an element g of $\overline{\mathrm{G}}(\mathbb{Q}_2)$ which is contained in the centralizer Z of a maximal split torus. A preimage $\tilde{g} \in \mathrm{Spin}_{f_1}$ of g under ϕ is given by:

$$\tilde{g} \quad = \quad \frac{2+\sqrt{2}}{4}\left(1+(\sqrt{2}-1)y_{r-1}x_{r-1}\right)\left(1+(\sqrt{2}-1)y_r x_r\right)\prod_{j=1}^{r-2}\left(\sqrt[4]{2}x_j y_j + \frac{1}{\sqrt[4]{2}}y_j x_j\right).$$

A direct computation shows that the cocycle $a \in \mathrm{H}^1(\mathbb{Q}_2, \mathrm{C})$ given by $a_\sigma = {}^\sigma\tilde{g}\tilde{g}^{-1}$ corresponds exactly to the element $2 \in \mathbb{Q}_2^*/(\mathbb{Q}_2^*)^4 \simeq \mathrm{H}^1(\mathbb{Q}_2, \mathrm{C})$. We check that $g \notin Z_c Z_s$ (see notation in [31, Sec. 2.5]) and so $2 \in \mathrm{H}^1(\mathbb{Q}_2, \mathrm{C})$ acts non-trivially on $\Delta_{(2)}$. Since this local Dynkin diagram has only one non-trivial symmetry, this implies $q' = 2$ (in case when $r = 2m+1$ with m odd).

For $r = 2m$ the type of G_1 is ${}^2\mathrm{D}_{2m}$ and the splitting field $\ell = \ell_1$ is the field $\mathbb{Q}[\sqrt{-3}]$. We have $h_\ell = 1$. By Proposition 4.7, $\mathbf{A} = \ell^*$. By Section 4.11, we obtain $\mathbf{A}_2/(\ell^*)^2 = \{\pm 1\}$. Again we can compute the value of q: the element $\sqrt{-3}$ represents a non-trivial element of $A_{\xi,R}$ which is not in $\mathbf{A}_2/(\ell^*)^2$, hence $q = 2$. Since $\pm 1, \pm\sqrt{-3}$ are the four elements composing $\ell^*_{(3)}/\left(\ell^*_{(3)}\right)^2 = \mathrm{H}^1(\mathbb{Q}_3, \mathrm{C})$, we see that $A_{\xi,R} \to \Xi_{(3)}$ is onto. This implies that $q' = \#\Xi_{(3)} = 2$.

Similar to Section 5.1, we have $[\Gamma_1 : \Lambda_1] = (q/q') \cdot \#\left(U_{\mathbf{A}}/U_\ell^{\mathfrak{n}}\right)$, where now $U_{\mathbf{A}}/U_\ell^{\mathfrak{n}}$ has order 2 if r is even and is trivial when r is odd. Hence from the above considerations we get:

$$[\Gamma_1 : \Lambda_1] \quad = \quad \begin{cases} 1 & \text{if } r = 2m+1 \text{ with } m \text{ even}, \\ 2 & \text{otherwise.} \end{cases} \tag{26}$$

6. UNIQUENESS OF Γ_0 AND Γ_1

The uniqueness part of Proposition 3.6 (resp. Proposition 3.9) implies that any principal arithmetic subgroup of H which has the same global type as Λ_0 (resp. Λ_1) must be commensurable with this latter group. More precisely, it shows that in each case there exists a uniquely determined group G/k and a collection of parahoric subgroups $(\mathrm{P}_v)_{v\in V_f}$, defined up to local conjugations, which are associated to our group Λ_i. In this section we will show that this defines Λ_i uniquely up to conjugation by the elements from $\overline{\mathrm{G}}(k)$.

6.1. Let $\mathscr{P} = (\mathrm{P}_v)_{v\in V_f}$ and $\mathscr{P}' = (\mathrm{P}'_v)_{v\in V_f}$ be two coherent collections of parahoric subgroups of G such that for all $v \in V_f$, P'_v is conjugate to P_v under an element of $\overline{\mathrm{G}}(k_v)$. For all but finitely many v, $\mathrm{P}_v = \mathrm{P}'_v$ and they are hyperspecial, hence there exists an element $g \in \overline{\mathrm{G}}(\mathbb{A}_f)$ such that $\mathscr{P}'$ is conjugate to $\mathscr{P}$ under g. Let $\overline{\mathrm{P}} = \prod_{v\in V_f} \overline{\mathrm{P}}_v$ be the stabilizer of $\mathscr{P}$ in $\overline{\mathrm{G}}(\mathbb{A}_f)$. The number of distinct $\overline{\mathrm{G}}(k)$-conjugacy classes of coherent collections $\mathscr{P}'$ as above is the cardinality $c(\overline{\mathrm{P}})$ of $\mathscr{C}(\overline{\mathrm{P}}) = \overline{\mathrm{G}}(k)\backslash\overline{\mathrm{G}}(\mathbb{A}_f)/\overline{\mathrm{P}}$, which is called the *class group* of $\overline{\mathrm{G}}$ relative to $\overline{\mathrm{P}}$. The *class number* $c(\overline{\mathrm{P}})$ is known to be finite (see e.g. [5, Prop. 3.9]). We need to compute its value for the groups constructed in Section 3.

We first consider a more general setup and then apply the results to our groups. Recall two isomorphisms (see [24, Prop. 8.8], a minor modification is needed in order to adjust the statement to our setting but the argument remains the same):

$$\begin{aligned}\overline{\mathrm{G}}(k)\backslash\overline{\mathrm{G}}(\mathbb{A}_f)/\overline{\mathrm{P}} &\simeq \overline{\mathrm{G}}(\mathbb{A}_f)/\overline{\mathrm{G}}(k)\cdot\overline{\mathrm{P}};\\ \overline{\mathrm{G}}(\mathbb{A}_f)/\overline{\mathrm{G}}(k)\cdot\overline{\mathrm{P}} &\simeq \delta_{\mathbb{A}_f}(\overline{\mathrm{G}}(\mathbb{A}_f))/\delta_{\mathbb{A}_f}(\overline{\mathrm{G}}(k)\cdot\overline{\mathrm{P}}),\end{aligned}$$

where $\delta_{\mathbb{A}_f}$ is the restriction of the product map $\prod_v \overline{\mathrm{G}}(k_v) \to \prod_v \mathrm{H}^1(k_v,\mathrm{C})$ to the finite adèle group $\overline{\mathrm{G}}(\mathbb{A}_f)$.

For every finite place v, $\mathrm{H}^1(k_v,\mathrm{G})$ is trivial (see [24, Theorem 6.4]) hence $\delta_v : \overline{\mathrm{G}}(k_v) \to \mathrm{H}^1(k_v,\mathrm{C})$ is surjective. Thus the image of $\delta_{\mathbb{A}_f}(\overline{\mathrm{G}}(\mathbb{A}_f))$ can be identified with the restricted direct product $\prod' \mathrm{H}^1(k_v,\mathrm{C})$ with respect to the subgroups $\delta_v(\overline{\mathrm{P}}_v)$. Also, $\delta_{\mathbb{A}_f}(\overline{\mathrm{G}}(k))$ identifies with the image of $\delta(\overline{\mathrm{G}}(k))$ under the natural map $\psi : \mathrm{H}^1(k,\mathrm{C}) \to \prod' \mathrm{H}^1(k_v,\mathrm{C})$. Hence we have an isomorphism

$$\delta_{\mathbb{A}_f}(\overline{\mathrm{G}}(\mathbb{A}_f))/\delta_{\mathbb{A}_f}(\overline{\mathrm{G}}(k)\cdot\overline{\mathrm{P}}) \simeq \textstyle\prod' \mathrm{H}^1(k_v,\mathrm{C})\big/\big((\psi\circ\delta)(\overline{\mathrm{G}}(k))\cdot\prod_v\delta_v(\overline{\mathrm{P}}_v)\big). \tag{27}$$

We can summarize the result as follows:

Proposition 6.2. *We have* $\mathscr{C}(\overline{\mathrm{P}}) \simeq \prod' \mathrm{H}^1(k_v,\mathrm{C})\big/\big((\psi\circ\delta)(\overline{\mathrm{G}}(k))\cdot\prod_v \delta_v(\overline{\mathrm{P}}_v)\big)$.

6.3. To apply this proposition to our groups we will describe $\prod' \mathrm{H}^1(k_v,\mathrm{C})$ and its quotient using idèle groups (see e.g. [15, Ch. VII, §3] for more about idèles). We first consider the case when G is not of type ${}^2\mathrm{D}_{2m+1}$. Then $\mathrm{H}^1(k_v,\mathrm{C})$ is canonically isomorphic to $(\ell\otimes_k k_v)^*/(\ell\otimes_k k_v)^{*\mathfrak{n}}$. Now, the group $\delta_v(\overline{\mathrm{P}}_v)$ corresponds to the kernel $\mathrm{H}^1(k_v,\mathrm{C})_{\xi_v}$ and by [5, Lemma 2.3 and Prop. 2.7] (see also Section 4.9), for all but a finite number of places (namely, $V_f \setminus (T_1 \cup R)$) this kernel is given by the elements having representatives in $\prod_{w|v}\mathfrak{o}_w^* \subset (\ell\otimes k_v)^*$, where $\mathfrak{o}_w^*$ is the group of integer units in ℓ_w. This shows that the restricted direct product $\prod' \mathrm{H}^1(k_v,\mathrm{C})$ is isomorphic to $J_f(\ell)/J_f(\ell)^{\mathfrak{n}}$, where $J_f(\ell)$ denotes the group of finite idèles of ℓ. We shall denote by $J_{\overline{\mathrm{P}}}/J_f(\ell)^{\mathfrak{n}}$ its subgroup corresponding to the product $\prod_v \delta_v(\overline{\mathrm{P}}_v)$ (where $J_{\overline{\mathrm{P}}} \subset J_f(\ell)$ contains $J_f(\ell)^{\mathfrak{n}}$). From the above explanation it follows that the group $J_{\overline{\mathrm{P}}}/J_f(\ell)^{\mathfrak{n}}$ differs only by a finite number of factors from the group $J_\circ(\ell)J_f(\ell)^{\mathfrak{n}}/J_f(\ell)^{\mathfrak{n}}$, where $J_\circ(\ell) < J_f(\ell)$ denotes the subgroup of finite idèles which are given by units at all places:

$$J_\circ(\ell) \quad = \quad \prod_{w\in V_f(\ell)} \mathfrak{o}_w^*.$$

We consider now the case ${}^2\mathrm{D}_{2m+1}$ which arises only for half of the groups G_0. In this case the group $\mathrm{H}^1(k_v,\mathrm{C})$ was described in Section 4.5 by the following exact sequence:

$$1 \to \mu_{\mathfrak{n}}(k_v)/N_{\ell|k}(\mathrm{R}_{\ell|k}(\mu_{\mathfrak{n}})(k_v)) \to \mathrm{H}^1(k_v,\mathrm{C}) \to (\ell\otimes_k k_v)^*/(\ell\otimes_k k_v)^{*\mathfrak{n}} \xrightarrow{N_{\ell|k}} k_v^*/k_v^{*\mathfrak{n}}.$$

We note that by [5, Sec. 5.3] together with the argument from Section 5.2, for every $v \in V_f$, the group $\mu_{\mathfrak{n}}(k_v)/N_{\ell|k}(\mathrm{R}_{\ell|k}(\mu_{\mathfrak{n}})(k_v))$ acts trivially on the local Dynkin diagram Δ_v and hence it is contained in $\delta_v(\overline{\mathrm{P}}_v)$. Therefore, using the same argument as above, we see that the quotient $\prod' \mathrm{H}^1(k_v,\mathrm{C})/\prod_v \delta_v(\overline{\mathrm{P}}_v)$ can be described as a quotient of $J_f(\mathbf{L})/J_f(\ell)^{\mathfrak{n}}$ by some subgroup $J_{\overline{\mathrm{P}}}/J_f(\ell)^{\mathfrak{n}}$ (with $J_f(\ell)^{\mathfrak{n}} \subset J_{\overline{\mathrm{P}}} \subset J_f(\mathbf{L})$). Here the idèle group $J_f(\mathbf{L})$ is defined by

$$J_f(\mathbf{L}) \quad = \quad \left\{(x_w)_w \in J_f(\ell) \mid N_{\ell|k}(x_w) \in (k_v^*)^4 \text{ for all } w|v,\ v\in V_f\right\},$$

and the group $\mathbf{L}$ introduced in Section 4.5 can be seen as a subgroup of $J_f(\mathbf{L})$. We can also consider $J_\circ(\mathbf{L}) = J_\circ(\ell)\cap J_f(\mathbf{L})$.

Recall that for the cases different from ${}^2\mathrm{D}_{2m+1}$ we have $\mathbf{L} = \ell^*$. We will unify the notation for idèles in the rest of the section and denote by $J_f(\mathbf{L})$ (resp. $J_\circ(\mathbf{L})$) the group $J_f(\ell)$ (resp. $J_\circ(\ell)$) if the group G is not of type ${}^2\mathrm{D}_{2m+1}$.

6.4. By Section 4.6, $\delta(\overline{\mathrm{G}}(k)) \supset A$, with the group A introduced there. By Proposition 4.7, in all the cases except when $k=\mathbb{Q}$ and G is of type ${}^2\mathrm{D}_{2m+1}$, the group A is a subgroup of $\ell^*/(\ell^*)^{\mathfrak{n}}$. By Proposition 3.9, when $k=\mathbb{Q}$ and $r = 2m+1$ our group G ($=\mathrm{G}_1$) is inner and hence the excluded case does not occur. So the group A can be identified with $\mathbf{A}/(\ell^*)^{\mathfrak{n}}$, where $\mathbf{A}$ is given by Proposition 4.7 *(1a), (1b), (2a)* or *(2c)*.

The group $J_{\overline{\mathrm{P}}}$ introduced in Section 6.3 can differ from $J_\circ(\mathbf{L})$ at the direct factors corresponding to the places of ℓ above the finite set $R\cup T_1$. But for groups G_0 (resp. G_1) under consideration, Section 5.2 (resp. Section 5.3) shows that we have an inclusion

$$\mathbf{A}J_\circ(\mathbf{L})J_f(\ell)^{\mathfrak{n}}/J_f(\ell)^{\mathfrak{n}} \quad \subset \quad \mathbf{A}J_{\overline{\mathrm{P}}}/J_f(\ell)^{\mathfrak{n}}. \tag{28}$$

It follows then from Proposition 6.2 and Section 6.3 that the class group $\mathscr{C}(\overline{\mathrm{P}})$ is a quotient of the group

$$\begin{aligned} \mathscr{C}_\circ &= (J_f(\mathbf{L})/J_f(\mathbf{L})^{\mathfrak{n}})/(\mathbf{A}J_\circ(\mathbf{L})J_f(\ell)^{\mathfrak{n}}/J_f(\ell)^{\mathfrak{n}}) \\ &\simeq J_f(\mathbf{L})/\mathbf{A}J_\circ(\mathbf{L})J_f(\ell)^{\mathfrak{n}}. \end{aligned} \tag{29}$$

6.5. To prove uniqueness it is sufficient to show that in each case under consideration the group $\mathscr{C}_\circ$ is trivial. We have the following commutative diagram with an exact row:

$$\begin{tikzcd} \mathbf{A} \arrow[r] \arrow[dr, "p_1"'] & J_f(\mathbf{L})/J_\circ(\mathbf{L})J_f(\ell)^{\mathfrak{n}} \arrow[r] & \mathscr{C}_\circ \arrow[r] & 1 \\ & \mathbf{L}/U_{\mathbf{L}} \arrow[u, "p_2"'] & & \end{tikzcd} \tag{30}$$

We need to show the surjectivity of the first horizontal map. To this end consider the composition $p_2 \circ p_1$. Since $h_\ell = 1$ the surjectivity of p_2 can be proved directly, expressing local uniformizers as the elements in ℓ. To prove the surjectivity of p_1 we must show that $\mathbf{A}U_{\mathbf{L}} = \mathbf{L}$. This is shown if $\mathbf{L}/\mathbf{A}$ has the same number of elements as $U_{\mathbf{L}}/U_{\mathbf{A}}$. With the description of $\mathbf{A}$ given in Proposition 4.7 and the information about $U_{\mathbf{A}}$ from Section 5.1 we can easily check this equality for all cases. This finishes the proof of uniqueness.

6.6. We showed that the groups Λ_0 and Λ_1 are defined uniquely up to conjugation in $\overline{H}$. It implies that the same is true for their normalizers Γ_0 and Γ_1. In the next three sections we will prove that these groups have minimal covolume among the corresponding lattices. The results of this section will then imply the uniqueness of the minimal orbifolds in Theorems 1 and 2.

7. PROOF OF THEOREM 1: ODD RANK

In this section we give a proof of Theorem 1 for the rank $r = 2m+1$. Here we assume that the minimal covolume lattice Γ is associated to an algebraic group G which is defined over $k \neq \mathbb{Q}$ and has the splitting field ℓ. By Section 2.3, G is of type ${}^2\mathrm{D}_{2m+1}$. Moreover, we can assume that $\Gamma = \Gamma^{\mathfrak{m}}$ (see Section 4.3). We would like to show that $\mathrm{G} = \mathrm{G}_0$. The results of Sections 3.7 and 4.3 would then imply immediately that the minimal covolume is attained on the lattice Γ_0.

7.1. Let us first consider the case $(k,\ell) = (k_0,\ell_0)$. Since the rational primes 2 and 3 are inert in k_0, the cardinality of the residue field $\mathfrak{f}_v$ is at least 4 for each $v \in V_f(k)$. By (5), this implies that $\lambda_v \geq 18$ for every possible lambda factor appearing in the covolume of Γ. Then if $T_1 \neq \emptyset$ we see that the covolume of Γ is strictly bigger than the covolume of Γ_0. Indeed, the index $[\Gamma : \Lambda]$ can differ from $[\Gamma_0 : \Lambda_0]$ only by the factors q and q' (the order of $\mathbf{A}_{\mathfrak{n}}/(\ell^*)^{\mathfrak{n}}$ depends only on ℓ), and by Proposition 4.10 we have $q \leq \prod_v \lambda_v$. Hence if $(k,\ell) = (k_0,\ell_0)$, we must have $T_1 = \emptyset$. By the uniqueness part of Proposition 3.9 we then have $\mathrm{G} = \mathrm{G}_0$.

Our discussion shows that the proof of the theorem reduces to proving the following statement: *If Γ is a cocompact arithmetic subgroup of the smallest covolume then the fields*

k, ℓ associated to it are the fields k_0, ℓ_0. The rest of this section is concerned with establishing this fact. The idea is to obtain bounds for the discriminants $\mathscr{D}_k$ and $\mathscr{D}_\ell$ from the inequality $\mu(H/\Gamma) \leq \mu(H/\Gamma_0)$ and the upper bound for $\mu(H/\Gamma_0)$, which we get from the estimates (7) and (8) (when $r \leq 16$) along with $[\Gamma_0 : \Lambda_0] = 2$.

7.2. From the volume formula and Proposition 4.12 it follows that for G of type ${}^2\mathrm{D}_{2m+1}$:

$$\mu(H/\Gamma) \geq \frac{1}{4^{\#T_1} h_\ell 2^{d+1}} \mathscr{D}_k^{r^2-r/2} \left(\frac{\mathscr{D}_\ell}{\mathscr{D}_k^2}\right)^{r-1/2} C(r)^d \mathscr{E}(\mathscr{P}),$$

where $\mathscr{P}$ is a coherent collection of parahoric subgroups defining a principal arithmetic subgroup whose normalizer is Γ. By [5, Appendix C], we have $4^{-\#T_1}\mathscr{E}(\mathscr{P}) > 1$ (this value corresponds to $\prod f_v$ in the latter reference), which gives:

$$\mu(H/\Gamma) > \frac{1}{h_\ell 2^{d+1}} \mathscr{D}_k^{r^2-r/2} \left(\frac{\mathscr{D}_\ell}{\mathscr{D}_k^2}\right)^{r-1/2} C(r)^d. \tag{31}$$

At this point we need to bound h_ℓ, and for this we will use the Brauer-Siegel theorem similarly to the way it is done in [5, proof of Prop. 6.1]. Since ℓ has signature $(2, d-1)$, the Brauer-Siegel theorem implies that for any $s > 1$:

$$h_\ell R_\ell \leq 2s(s-1)2^{-2}\Gamma\left(\frac{s}{2}\right)^2 \Gamma(s)^{d-1}(2^{-2d+2}\pi^{-2d}\mathscr{D}_\ell)^{s/2}\zeta(s)^{2d}, \tag{32}$$

where R_ℓ is the regulator of ℓ. In [10] it is proved that, with the exception of three fields, the regulator of a number field is greater than $1/4$. Since these three exceptional fields are totally complex, they cannot arise as ℓ. With $R_\ell \geq 1/4$ and taking $s = 2$ in (32), we get

$$h_\ell \leq 16\left(\frac{\pi}{12}\right)^{2d} \mathscr{D}_\ell. \tag{33}$$

Combining (31) and (33), and in the second step using $\mathscr{D}_\ell \geq \mathscr{D}_k^2$, we can write:

$$\mu(H/\Gamma) > \frac{1}{32} \mathscr{D}_k^{r^2-5r/2+1} \mathscr{D}_\ell^{r-3/2} a(r)^d \tag{34}$$

$$\geq \frac{1}{32} \mathscr{D}_k^{r^2-r/2-2} a(r)^d, \tag{35}$$

where

$$a(r) = \frac{1}{2}\left(\frac{12}{\pi}\right)^2 C(r).$$

7.3. We now reduce to a finite number of possibilities for the field k by bounding the discriminant $\mathscr{D}_k$. For $r \geq 15$ the factor $a(r)$ is greater than one, so its power to d can be replaced in (35) by a square without changing the inequality. Comparing this value with the upper bound for the covolume of Γ_0 we get:

$$\left(\frac{\mathscr{D}_k}{5 \cdot 11^{1/r}}\right)^{r^2-r/2-2} \leq 128\left(\frac{\pi}{12}\right)^4 5^2 \cdot 11^{2/r}, \tag{36}$$

which is only possible for $\mathscr{D}_k \le 6.1$ if $r \ge 15$ (the worst bound appears with the smallest r, here $r = 15$). This implies that $k = k_0$.

For $r = 3, 5, \ldots, 13$ the factor $a(r)$ is less than one and we have to be more careful in order to exclude fields k of high degrees. We use *unconditional* (i.e. not depending on the Generalized Riemann Hypothesis) bounds for discriminants of totally real number fields given in [20, table 4]. Taking $d \ge 7$ we have $\mathscr{D}_k > 9.3^d$, and we can check that for all $r = 3, 5, \ldots, 13$ the factor $9.3^{r^2-r/2-2}a(r)$ is greater than 1. From (34) we then obtain:

$$\mu(H/\Gamma) \quad > \quad \frac{1}{32}\left(9.3^{r^2-r/2-2}a(r)\right)^7, \tag{37}$$

and we check that for each $r = 3, 5, \ldots, 13$ the right hand side of (37) is greater than the covolume of Γ_0. So the degree of k must be strictly less than 7. For the degrees $d = 2, \ldots, 6$ we compare the bound (35) with the covolume of Γ_0. In each degree and for each r it gives a bound for $\mathscr{D}_k$. We list the bounds we get for $r = 3$, which are worst than for other r:

$$\begin{aligned} d = 2 \quad &: \quad \mathscr{D}_k \le 22;\\ d = 3 \quad &: \quad \mathscr{D}_k \le 198;\\ d = 4 \quad &: \quad \mathscr{D}_k \le 1\,778;\\ d = 5 \quad &: \quad \mathscr{D}_k \le 15\,956;\\ d = 6 \quad &: \quad \mathscr{D}_k \le 143\,195. \end{aligned}$$

The bounds for $\mathscr{D}_k$ are sufficient to determine from the tables of totally real number fields (see [32] or [33]) precisely which fields may occur as k. For $r = 7, 9, 11, 15$ we see immediately that the only possibility is $k = k_0$. For $r = 5$ the quadratic field of discriminant 8 is not excluded, while for $r = 3$ we are left with fourteen possible fields, all of degree lower than 5.

7.4. We now turn to the problem of excluding all but a finite number of extensions $\ell|k$. For $r \ge 15$ and odd we proved that $\mathscr{D}_k = 5$ and from (34) we can give a bound for $\mathscr{D}_\ell$ similar to the way we obtained the bound (36) for $\mathscr{D}_k$. We obtain that $\mathscr{D}_\ell \le 336$. There are three such field of signature $(2,1)$, but only $\ell = \ell_0$ is an extension of k_0.

For $r = 3$ the only possible field of degree $d = 5$ is k with $\mathscr{D}_k = 14\,641$. Comparing the bound (34) for this given value of $\mathscr{D}_k$ with $\mu(H/\Gamma_0)$, we get $\mathscr{D}_\ell \le 731 \cdot 10^6$. The degree 10 is too large to find a list of possible number fields in tables from [32] and [33]. But if we make use of the information that ℓ is an extension of a totally real quintic field, we can use the results of [27] where the smallest discriminants of such fields are determined for each signature. It turns out that our bound for $\mathscr{D}_\ell$ is smaller than the smallest possible value of the discriminant ($= 1\,332\,031\,009$). This implies that no extension of k satisfies the inequality, and hence the quintic field is eliminated. For $d = 4$ we have three possibilities for k with $\mathscr{D}_k = 725$, $1\,125$ and $1\,600$. From [8] we known that here $\mathscr{D}_\ell \ge 4\,286\,875$. Similarly to what was done for the quintic field above, we can exclude the case $\mathscr{D}_k = 1\,600$. For the others two fields this argument does not work. However, with the method based on the class field theory which is explained in [8], we can (using e.g. Pari/GP) find all quadratic extensions of k with real signature 2 up to the bound for $\mathscr{D}_\ell$. It turns out that for $\mathscr{D}_k = 1\,125$ there are no such extensions. For $\mathscr{D}_k = 725$ two extensions ℓ are possible (with discriminants $\mathscr{D}_\ell = 5\,781\,875$ and $9\,986\,875$, resp.), both with class number $h_\ell = 1$. With this information at hand we can come back to the inequality (31) and refine our bound for $\mathscr{D}_k$. This gives us the bound $\mathscr{D}_k \leq 699$, which excludes the last quartic case $\mathscr{D}_k = 725$. For $d = 2$, resp. $d = 3$, the maximal value for $\mathscr{D}_\ell$ arises when $\mathscr{D}_k = 5$, resp. $\mathscr{D}_k = 49$. Using this value of $\mathscr{D}_k$ in (34) and comparing with $\mu(H/\Gamma_0)$ we get the following bounds for $\mathscr{D}_\ell$:

$$\begin{aligned} d = 2 \quad &: \quad \mathscr{D}_\ell \leq 5\,893; \\ d = 3 \quad &: \quad \mathscr{D}_\ell \leq 409\,830. \end{aligned}$$

We can check in the tables from [32] and [33] that all fields ℓ respecting these bounds have class number one. As above, this allows us to refine the bound for $\mathscr{D}_k$ by putting $h_\ell = 1$ in (31). We find that the only possible values of $\mathscr{D}_k$ are $5, 8, 12$ for $d = 2$ and 49 for $d = 3$. For each of these $\mathscr{D}_k$ we now refine the bound for $\mathscr{D}_\ell$ using $h_\ell = 1$. It turns out that only $k = k_0$ has possible extensions ℓ whose discriminant is small enough, and these possible extensions are given by ℓ with $\mathscr{D}_\ell = 275$, 400 and 475.

For $r = 5$ the bound for $\mathscr{D}_\ell$ in case $\mathscr{D}_k = 8$ is small enough to exclude this field. For $r = 5, \ldots, 13$ we then have necessarily $\mathscr{D}_k = 5$, for which the bounds for $\mathscr{D}_\ell$ allow us to check in a first step that $h_\ell = 1$. Again this improves the bounds for $\mathscr{D}_\ell$. From this we find that ℓ must be ℓ_0.

7.5. To achieve the goal of this section it remains to exclude the cases $\mathscr{D}_\ell = 400$ or 475 when $r = 3$. These two discriminants correspond to the extensions $k[\sqrt{\omega}]$ and $k[\sqrt{\beta}]$ of $k = k_0$, where

$$\omega = \frac{1+\sqrt{5}}{2} \quad \text{and} \quad \beta = -1 + 2\sqrt{5}.$$

The fundamental units in $k[\sqrt{\omega}]$ (resp. in $k[\sqrt{\beta}]$) are given by $\tau_1 = \sqrt{\omega}$ and $\tau_2 = 1 + \sqrt{\omega}$ (resp. $\tau_1 = \frac{1+\sqrt{\beta}}{2}$ and $\tau_2 = \frac{1-\sqrt{\beta}}{2}$). We can compute for these two fields ℓ that

$$\#(\mathbf{A}_4/(\ell^*)^4) = 4.$$

This computation is carried out the same way as it was done for ℓ_0 in Section 5.1. This gives $[\Gamma : \Lambda] \leq 4^{1+\#T_1}$, and with this new bound we can check that the covolume of Γ is bigger than $\mu(H/\Gamma_0)$.

We showed that for $r = 2m+1$ the minimal covolume in the cocompact case is attained on the group Γ_0 constructed in Section 3. Together with the index computation in Section 5 and the uniqueness result of Section 6 it finishes the proof of Theorem 1 for odd rank.

8. PROOF OF THEOREM 1: EVEN RANK

8.1. We now deal with the even rank case in Theorem 1. Admissible groups are of type ${}^2\mathrm{D}_{2m}$ or ${}^{3,6}\mathrm{D}_4$. The proof follows the same idea as in odd rank and we will allow ourselves to skip some details which are clear from Section 7. Thus, repeating the argument from

Section 7.1 we obtain that it is enough to show that the minimal covolume subgroups are attached to $(k,\ell) = (k_0,\ell_0)$. In particular, in rank 4 we need to exclude the triality forms ${}^{3,6}\mathrm{D}_4$. As in Section 7, we denote by $\Gamma = \Gamma^{\mathrm{m}}$ a subgroup of G defined over $k \neq \mathbb{Q}$ with a splitting field ℓ and we suppose that Γ has minimal covolume. We will treat the case ${}^{3,6}\mathrm{D}_4$ separately in Section 8.5 and assume in the beginning that G is of type ${}^2\mathrm{D}_{2m}$.

8.2. The index $[\Gamma : \Lambda]$ which appears in the formula for $\mu(H/\Gamma)$ is bounded by the integer $2^{2d-1}2^{\#R}4^{\#T_1}h_\ell$ (by Proposition 4.12). The factor $4^{\#T_1}$ can be eliminated the same way as it was done in Section 8.2. From [25, Appendix] we have the inequality

$$2^{\#R} \leq \frac{\mathscr{D}_\ell}{\mathscr{D}_k^{[\ell:k]}}. \tag{38}$$

The bound (33) for h_ℓ is still valid in the current setting. Combining all these facts together we get

$$\mu(H/\Gamma) > \frac{1}{8}\mathscr{D}_k^{r^2-5r/2+3}\mathscr{D}_l^{r-5/2}a(r)^d \tag{39}$$

$$\geq \frac{1}{8}\mathscr{D}_k^{r^2-r/2-2}a(r)^d, \tag{40}$$

where now

$$a(r) = \frac{1}{4}\left(\frac{12}{\pi}\right)^2 C(r).$$

8.3. For $r \geq 16$ we can check that $a(r) \geq 1$ and this allows us to bound $\mathscr{D}_k$ by an inequality similar to (36). We get $\mathscr{D}_k \leq 5.9$, proving that $k = k_0$ if $r \geq 16$. For $r = 4,\ldots,14$ we exclude the fields k of degree $d \geq 5$ using the same trick as in Section 7.3 (Odlyzko's bound employed here is $\mathscr{D}_k \geq (6.5)^d$ for $d \geq 5$). Comparing (40) with $\mu(H/\Gamma_0)$ we get bounds for $\mathscr{D}_k$ depending on d for each $r = 4,\ldots,14$. In particular, for $r = 4$:

$$\begin{aligned} d = 2 &: \mathscr{D}_k \leq 12; \\ d = 3 &: \mathscr{D}_k \leq 62; \\ d = 4 &: \mathscr{D}_k \leq 323. \end{aligned}$$

We see that k can be different from k_0 only when $r = 4$ (with possible quadratic fields of discriminants $\mathscr{D}_k = 8, 12$ and a cubic field of discriminant $\mathscr{D}_k = 49$), or when $r = 6$ (with $\mathscr{D}_k = 8$).

8.4. We now check possibilities for the field ℓ. For $r \geq 16$ we obtain the bound $\mathscr{D}_\ell \leq 479$. Besides the field ℓ_0, we then have a possibility of ℓ having discriminant 400 or 475. These fields have class number $h_\ell = 1$ which allows us to get an inequality

$$\left(\frac{\mathscr{D}_\ell}{25\cdot 11}\right)^{r-3/2} \leq 8\cdot 11.$$

From this inequality we obtain that for $r \geq 16$, $\mathscr{D}_\ell \leq 383$, which implies that $\ell = \ell_0$.

In rank $r = 4$, for the only possible cubic field k with $\mathscr{D}_k = 49$ we obtain $\mathscr{D}_\ell \leq 16\,809$, and no any field of signature $(4,1)$ satisfies this condition. Similarly, the cases $\mathscr{D}_k = 12$ for $r = 4$ and $\mathscr{D}_k = 8$ for $r = 6$ can be also eliminated. Therefore, we obtain the bound $\mathscr{D}_\ell \leq 28\,662$ which is valid in rank $r = 4$ (with the worst case corresponding to $\mathscr{D}_k = 5$) as well as in higher ranks. We can check in the table from [33] that admissible quartic fields ℓ whose discriminant respects this bound have class number $h_\ell \leq 3$. This information can be used in the inequality

$$\mu(H/\Gamma) \quad \geq \quad \frac{1}{2^{2d-1}h_\ell}\mathscr{D}_k^{r^2-5r/2+3}\mathscr{D}_\ell^{r-3/2}C(r)^d, \tag{41}$$

to get the refinement $\mathscr{D}_\ell \leq 2\,064$. All fields ℓ which satisfy this condition have the class number $h_\ell = 1$. Again, this improves the bound for the discriminant to $\mathscr{D}_\ell \leq 1\,330$ when $r = 4$ and $\mathscr{D}_k = 5$. For $\mathscr{D}_k = 8$ and the same rank we have a better bound $\mathscr{D}_\ell \leq 224$ which excludes this case and proves that $k = k_0$ also in rank $r = 4$. For the field ℓ we now have the possibilities $\mathscr{D}_\ell = 275, 400, 475, 775$. For each of these fields the extension $\ell|k_0$ contains exactly one ramified place, which means that $\#R = 1$. The inequality (38) can be advantageously replaced by $2^{\#R} = 2$. This gives:

$$\mu(H/\Gamma) \quad \geq \quad \frac{1}{2^3}5^{r^2-r/2}\left(\frac{\mathscr{D}_\ell}{5^2}\right)^{r-1/2}C(r)^2. \tag{42}$$

Comparing the value of this bound with $\mu(H/\Gamma_0)$ we can check that $\ell = \ell_0$, with the only possible exception in rank 4 of the field ℓ with $\mathscr{D}_\ell = 400$. Now for this specific ℓ we can compute that $\#\mathbf{A}_2/(\ell^*)^2 = 2$ (we use the description of ℓ given in Section 7.5). Then we have $[\Gamma : \Lambda] \leq 4^{1+\#T_1}$ which allows us to show that $\mu(H/\Gamma)$ is bigger than $\mu(H/\Gamma_0)$ in the case $\mathscr{D}_\ell = 400$. Thus, in rank $r = 4$ we have $\ell = \ell_0$ as well.

8.5. To complete the proof of Theorem 1 we need to exclude the arithmetic 7-orbifolds defined by triality forms. So assume now that G is of type ${}^{3,6}\mathrm{D}_4$. In this case we have $[\ell : k] = 3$ and $[\Gamma : \Lambda] \leq 2^{3d+1}2^{\#R}4^{\#T_1}h_\ell$. Moreover, the bound (33) should be replaced by $h_\ell \leq 16(\pi/12)^{3d}\mathscr{D}_\ell$. It follows that

$$\mu(H/\Gamma) \quad \geq \quad \frac{1}{32}a^d\mathscr{D}_k^{11},$$

where $a = 1/8(12/\pi)^3C(4)$. For $d \geq 5$ we can check using $\mathscr{D}_k \geq 6.5^d$ that $\mu(H/\Gamma)$ is necessarily bigger than $\mu(H/\Gamma_0)$ and for smaller d we have:

$$\begin{aligned} d = 2 &\quad : \quad \mathscr{D}_k \leq 15; \\ d = 3 &\quad : \quad \mathscr{D}_k \leq 86; \\ d = 4 &\quad : \quad \mathscr{D}_k \leq 490. \end{aligned}$$

In particular, the totally real field k cannot be of degree $d = 4$. The inequality

$$\mu(H/\Gamma) \quad \geq \quad \frac{1}{32}a^d\mathscr{D}_k^{13/2}\mathscr{D}_\ell^{3/2},$$

gives us different bounds for $\mathscr{D}_\ell$ for each of the possible $\mathscr{D}_k$. For $\mathscr{D}_k = 5$, $d = 2$ and $\mathscr{D}_k = 49$, $d = 3$ we have the worst possible bounds:

$$\begin{aligned} d = 2 \quad &: \quad \mathscr{D}_\ell \leq 445\,619; \\ d = 3 \quad &: \quad \mathscr{D}_\ell \leq 7.7 \cdot 10^6. \end{aligned}$$

We can check in the table from [33] that all fields ℓ of degree 6 with at least one real place (condition (3) from Proposition 2.5) and with $\mathscr{D}_\ell \leq 445\,619$ have the class number $h_\ell = 1$. This information allows us to improve the bound to $\mathscr{D}_\ell \leq 20\,165$, and no any admissible field ℓ has the discriminant small enough. Thus the case $d = 2$ is excluded. For $d = 3$ the bound we have is in contradiction with Odlyzko's bound $\mathscr{D}_\ell > (6.1)^9$, given in [21].

This finishes the proof of Theorem 1.

9. PROOF OF THEOREM 2

9.1. We begin the proof of Theorem 2 with the easiest case $r = 2m+1$ with m even. The same argument as in Section 7.1 implies that it is enough to show that a minimal covolume arithmetic subgroup $\Gamma = \Gamma^{\mathrm{m}}$ is necessarily defined by a group G of inner type. Thus it suffices to show that $\mathscr{D}_\ell = 1$. From Proposition 4.12, the bound (33) for h_ℓ and the inequality $4^{-\#T_1}\mathscr{E}(\mathscr{P}) > 1$, we get

$$\mu(H/\Gamma) \quad \geq \quad \frac{1}{8 \cdot 16}\left(\frac{12}{\pi}\right)^2 \mathscr{D}_\ell^{r-3/2}.$$

Since $r \geq 5$ this gives the bound $\mathscr{D}_\ell \leq 2.27$, proving that $\ell = \ell_1 = \mathbb{Q}$.

9.2. Let us assume now that $r = 2m+1$ with m odd. Since the result of Theorem 2 for $r = 3$ follows from [13], we allow ourselves to present here the proof with the assumption that $r \neq 3$. The proof in the case $r = 3$ with our method is also possible but it requires a bit more effort (see [9, §15.9]).

So we have $r \geq 7$. We can check that the product

$$\zeta(2)\zeta(4)\cdots\zeta(2r-2)\zeta(r)$$

appearing in $\mu(H/\Lambda_1)$ is bounded above by 2. Moreover, the lambda factor $\lambda_{(2)}$ can be bounded by 2^{2r-1}. Thus dividing $\mu(H/\Lambda_1)$ by the index $[\Gamma_1 : \Lambda_1]$ from (26), we obtain:

$$\mu(H/\Gamma_1) \quad < \quad \frac{4^{r-1/2}}{3} C(r).$$

We can compare this value with the general bound for a subgroup $\Gamma = \Gamma^{\mathrm{m}}$ obtained by our usual argument (Proposition 4.12, bound for h_ℓ, etc.):

$$\mu(H/\Gamma) \quad \geq \quad \frac{1}{8 \cdot 16}\left(\frac{12}{\pi}\right)^2 \mathscr{D}_\ell^{r-3/2}.$$

For $r \geq 7$, if we suppose that Γ is of the smallest covolume, we must have $\mathscr{D}_\ell \leq 6.3$. So $\mathscr{D}_\ell = 1$ and $\mathscr{D}_\ell = 5$ are the only possibilities (recall that by Proposition 2.5, ℓ must be totally real). In particular, $h_\ell = 1$ and this gives a better bound for $\mu(H/\Gamma)$, which is enough to get $\mathscr{D}_\ell \leq 4.7$. The latter is satisfied only for $\ell = \mathbb{Q}$. Hence the group Γ of minimal covolume must be associated to a group G of inner type ${}^1\mathrm{D}_{2m+1}$. Thus G is inner over each place of $\mathbb{Q}$. Moreover, we know that $G = \mathrm{Spin}_f$ for some quadratic form f over $\mathbb{Q}$. But since over $\mathbb{R}$ the Hasse symbol of f must by the admissibility condition be equal to -1, the Hilbert's reciprocity [22, Ch. VII] implies the existence of at least one finite place $v = (p)$ where G is non-split. This place v appears in the formula for the covolume of Γ with a non-trivial lambda factor and we can check using (9) that in case $v \neq (2)$ we have $\lambda_v/4 \geq \lambda_{(2)}$. Thus considering all other non-trivial λ_v, we get $\prod_v \lambda_v \cdot 4^{-\#T_1} \geq \lambda_{(2)}$. This implies that the smallest volume is obtained when G is non-split only at $v = (2)$. This is exactly the case for $\mathrm{G} = \mathrm{G}_1$, proving the minimality of the covolume of Γ_1.

9.3. We finally deal with the last case $r = 2m$. Again, even though the proof for $r = 4$ is possible, we will assume that $r \geq 6$ and refer to [13] for the remaining case. In the covolume of Γ_1 we can bound the Euler product by 2. Let us consider the lattice Γ of the minimal covolume. By the admissibility condition, the group G defining Γ must be of outer type ${}^2\mathrm{D}_{2m}$. Comparing $\mu(H/\Gamma_1)$ with $\mu(H/\Gamma)$ we have:

$$3^{r-1/2} \quad > \quad \frac{1}{4 \cdot 2^{\#R} h_\ell} \mathscr{D}_\ell^{r-1/2}. \tag{43}$$

We use the inequality $2^{\#R} \leq \mathscr{D}_\ell$ to simplify the right hand side. We can use the bound (33) which is valid here but requires a different argument: Indeed, this bound was obtained under the assumption that ℓ contains only two roots of unity. In the current setting it is no longer the case, since e.g. $\ell = \mathbb{Q}[i]$ has 4 roots of unity. But this is the only exception and $\mathbb{Q}[i]$ has the class number 1 and hence satisfies the bound. Thus for $r \geq 6$ we obtain $\mathscr{D}_\ell \leq 8$. It follows that $\#R = 1$ and $h_\ell = 1$. Coming back to (43) with the new information at hand, we get $\mathscr{D}_\ell \leq 4$. To exclude the case $\mathscr{D}_\ell = 4$, we compute the order of $\mathbf{A}_2/(\ell^*)^2$ for $\ell = \mathbb{Q}[i]$. By Proposition 4.7, we have $\mathbf{A} = \ell$ and using the method of Section 4.11 we obtain easily that $\ell_2/(\ell^*)^2$ has order 2. Compared to the general bound for the index in Proposition 4.12, this improves the bound by a factor two: $[\Gamma : \Lambda] \leq 2 \cdot 2 \cdot 4^{\#T_1}$. With this new bound for the index we can show that $\mu(H/\Gamma)$ is bigger than $\mu(H/\Gamma_1)$. It follows that the group Γ of the smallest covolume is necessarily associated to a group G with the splitting field $\ell = \ell_1$. We can see that the smallest covolume is obtained when no lambda factor appears in the volume formula. This implies that $\mathrm{G} = \mathrm{G}_1$ and thus completes the proof of Theorem 2.

10. GROWTH OF THE MINIMAL VOLUME AND PROOF OF THEOREM 3

10.1. The formulas from Theorems 1 and 2 allow us to investigate the behavior of the minimal volume as a function of the dimension of the space. As before, we will treat the compact and non-compact cases separately.

By Theorem 1 we have

$$\mathrm{Vol}_{\mathbb{H}}(O_0^n) = \frac{5^{r^2-r/2}\cdot 11^{r-1/2}\cdot(r-1)!}{2^{2r-1}\pi^r} L_{\ell_0|k_0}(r)\prod_{i=1}^{r-1}\frac{(2i-1)!^2}{(2\pi)^{4i}}\zeta_{k_0}(2i),$$

where $r=\frac{n+1}{2}$. The product $Z = L_{\ell_0|k_0}(r)\zeta_{k_0}(2)\cdots\zeta_{k_0}(2r-2)$ is minimal for $r=3$, and by evaluation we get $Z > 1.14$. Thus we have the following lower bound

$$\mathrm{Vol}_{\mathbb{H}}(O_0^n) > 1.14\cdot\frac{5^{r^2-r/2}\cdot 11^{r-1/2}\cdot(r-1)!}{2^{2r-1}\pi^r}\prod_{i=1}^{r-1}\frac{(2i-1)!^2}{(2\pi)^{4i}}. \tag{44}$$

It is easy to see that for $r\geq 15$, the product $\prod_{i=1}^{r-1}\frac{(2i-1)!^2}{(2\pi)^{4i}} > 1$. This leads to a lower bound

$$\mathrm{Vol}_{\mathbb{H}}(O_0^n) > (r-1)! \quad \text{for } r\geq 15,$$

which implies that the minimal volume in Theorem 1 *grows super-exponentially* for high enough n.

Similar argument works for the volumes of the non-compact orbifolds from Theorem 2. Here we have

$$\mathrm{Vol}_{\mathbb{H}}(O_1^n) = C_1(r)\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}}\zeta(2i) > 1.78\cdot C_1(r)\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}}, \tag{45}$$

where $r=\frac{n+1}{2}$ and $C_1(r)$ depends on $r \pmod 4$. We can show that for $r\geq 20$, the product $\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}} > (2(r-1)-1)!$ and $C_1(r) > 1$ if $r\not\equiv 1 \pmod 4$, $C_1(r) > \frac{1}{2^{r-2}}$, otherwise. Hence in all the cases

$$\mathrm{Vol}_{\mathbb{H}}(O_1^n) > r! \quad \text{for } r\geq 20.$$

Again we see that the minimal volume grows super-exponentially for high enough n.

10.2. We can compute the actual values of the minimal volume for small dimensions n. Table 2 shows the results of a numerical computation which was done using Pari/GP calculator. It gives *approximate* values of the minimal volume for each of the cases.

Our computation and estimation show that the minimal volume decrease with n till $n=7$ (resp. $n=17$) in the compact case (resp. non-compact case). After this it starts to grow eventually reaching a very fast super-exponential growth which we expect from the estimate in Section 10.1.

10.3. We are now ready to prove Theorem 3 from the introduction. The ratio $Q(n) = \mathrm{Vol}_{\mathbb{H}}(O_0^n)/\mathrm{Vol}_{\mathbb{H}}(O_1^n)$ can be bounded from below as follows.

Similar to the previous considerations we can show that for odd n,

$$\mathrm{Vol}_{\mathbb{H}}(O_1^n) \leq \frac{4^{r-1/2}}{2^{r-1}}2\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}} = 2^{r+1}\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}}.$$

n	5	7	9	11	13	15	17	19	21	23	25	27	29
$\mathrm{Vol}_{\mathbb{H}}(O_0^n)$	$1.53\cdot 10^{-3}$	$5.45\cdot 10^{-4}$	$7.20\cdot 10^{-3}$	9.87	$3.14\cdot 10^{6}$	$4.57\cdot 10^{14}$	$5.36\cdot 10^{25}$	$8.40\cdot 10^{39}$	$2.75\cdot 10^{57}$	$2.79\cdot 10^{78}$	$1.27\cdot 10^{103}$	$3.62\cdot 10^{131}$	$8.79\cdot 10^{163}$
$\mathrm{Vol}_{\mathbb{H}}(O_1^n)$	$3.65\cdot 10^{-4}$	$1.89\cdot 10^{-6}$	$9.30\cdot 10^{-11}$	$3.52\cdot 10^{-11}$	$1.20\cdot 10^{-12}$	$3.52\cdot 10^{-14}$	$2.07\cdot 10^{-18}$	$2.69\cdot 10^{-14}$	$3.65\cdot 10^{-12}$	$1.13\cdot 10^{-10}$	$6.63\cdot 10^{-13}$	$1.25\cdot 10^{-2}$	$1.98\cdot 10^{5}$

TABLE 2. Approximate volumes of O_0^n and O_1^n for small n

It implies

$$Q(n) \geq \frac{5^{r^2-r/2}\cdot 11^{r-1/2}\cdot (r-1)!}{2^{2r-1}\pi^r\cdot 2^{r+1}}\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}}.$$

Now as in Section 10.1, we have $\prod_{i=1}^{r-1}\frac{(2i-1)!}{(2\pi)^{2i}} > 1$ for $r \geq 15$, which implies that for odd $n \geq 29$, we have $Q(n) > (r-1)!$ where $r = \frac{n+1}{2}$. Together with the computational data from Table 2 this finishes the proof of Theorem 3 for odd n. The case of even dimension is obtained by an entirely similar application of the results from [2].

REFERENCES

[1] I. Agol, M. Belolipetsky, P. Storm, K. Whyte, Finiteness of arithmetic hyperbolic reflection groups, *Groups, Geometry, and Dynamics*, **2** (2008), 481–498.

[2] M. Belolipetsky, On volumes of arithmetic quotients of SO$(1,n)$, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5)*, **3** (2004), 749–770; Addendum: *ibid.*, **6** (2007), 263–268.

[3] A. Borel, Commensurability classes and volumes of hyperbolic 3-manifolds, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, **8** (1981), 1–33.

[4] A. Borel, Linear algebraic groups (2nd enlarged ed.), *Graduate Texts in Math.*, **126**, Springer (1991).

[5] A. Borel, G. Prasad, Finiteness theorems for discrete subgroups of bounded covolume in semi-simple groups, *Inst. Hautes Études Sci. Publ. Math.*, **69** (1989), 119–171; Addendum: *ibid.*, **71** (1990), 173–177.

[6] T. Chinburg, E. Friedman, The smallest arithmetic hyperbolic three-orbifold, *Invent. Math.*, **86** (1986), 507–527.

[7] V. I. Chernousov, A. A. Ryzhkov, On the classification of maximal arithmetic subgroups of simply connected groups, *Sb. Math.*, **188** (1997), 1385–1413.

[8] H. Cohen, F. Diaz y Diaz, M. Olivier, Computing ray class groups, conductors and discriminants, *Math. Comp.*, **67** (1998), 773–795.

[9] V. Emery, Du volume des quotients arithmétiques de l'espace hyperbolique, *PhD thesis*, Université de Fribourg (2009).

[10] E. Friedman, Analytic formulas for the regulator of a number field, *Invent. Math.*, **98** (1989), no. 3, 599–622.

[11] M.-A. Knus, A. Merkujev, M. Rost, J.-P. Tignol, The book of involutions, *Colloquium Publications.*, **44**, American Mathematical Society (1998).

[12] S. Helgason, Differential geometry and symmetric spaces, *Pure and Appl. Math.*, **12**, Academic Press (1962).

[13] T. Hild, The cusped hyperbolic orbifolds of minimal volume in dimensions less than ten, *J. Algebra*, **313** (2007), 208–222.

[14] T. Hild, R. Kellerhals, The FCC lattice and the cusped hyperbolic 4-orbifold of minimal volume: In memoriam H. S. M. Coxeter, *J. Lond. Math. Soc., II. Ser.*, **75** (2007) No. 3, 677–689.

[15] S. Lang, Algebraic number theory, *Graduate Texts in Math.*, **110**, Springer (1994).

[16] J.-S. Li, J. Millson, On the first Betti number of a hyperbolic manifold with an arithmetic fundamental group, *Duke Math. J.*, **71** (1993), 365–401.

[17] C. Maclachlan, A. W. Reid, The arithmetic of hyperbolic 3-manifolds, *Graduate Texts in Math.*, **219**, Springer (2003).

[18] R. Meyerhoff, The cusped hyperbolic 3-orbifold of minimum volume, *Bull. Am. Math. Soc., New Ser.*, **13** (1985), 154–156.

[19] A. Mohammadi, A. Salehi Golsefidy, Discrete subgroups acting transitively on vertices of a Bruhat-Tits building, *Duke Math. J.*, to appear.

[20] A. M. Odlyzko, Bounds for discriminants and related estimates for class numbers, regulators and zeros of zeta functions: A survey of recent results, *Sémin. Théor. Nombres Bordeaux, Sér. II*, **2** (1990), 119–141.

[21] A. M. Odlyzko, Discriminant bounds, http://www.dtc.umn.edu/∼odlyzko/unpublished/index.html

[22] O. O'Meara, Introduction to quadratic forms, *Grundl. der Math. Wiss.*, **117**, Springer (1963).

[23] T. Ono, On algebraic groups and discontinuous groups, *Nagoya Math. J.*, **27** (1966), 279–322.

[24] V. P. Platonov, A. S. Rapinchuk, Algebraic groups and number theory (English translation), *Pure and Appl. Math*, **139**, Academic Press (1994).

[25] G. Prasad, Volumes of S-arithmetic quotients of semi-simple groups, *Inst. Hautes Études Sci. Publ. Math.*, **69** (1989), 91–117.

[26] J. Rohlfs, Die maximalen arithmetisch definierten Untergruppen zerfallender einfacher Gruppen, *Math. Ann.*, **244** (1979), 219–231.

[27] S. Selmane, Quadratic extensions of totally real quintic fields, *Math. Comp.*, **70** (2001), 837–843.

[28] C. L. Siegel, Some remarks on discontinuous groups, *Ann. of Math.*, **46** (1945), 708–718.

[29] W. Scharlau, Quadratic and Hermitian forms, *Grundl. der Math. Wiss.*, **270**, Springer (1985).

[30] J. Tits, Classification of algebraic semisimple groups, *Proc. Symp. Pure Math.*, **9** (1966), 33–62.

[31] J. Tits, Reductive groups over local fields, *Proc. Symp. Pure Math.*, **33** (1979), Part I, 29–69.

[32] The Bordeaux Database, ftp://megrez.math.u-bordeaux.fr/pub/numberfields/

[33] QaoS online database, http://qaos.math.tu-berlin.de/cgi-bin/qaos/query.scm

IMPA, ESTRADA DONA CASTORINA, 110, 22460-320 RIO DE JANEIRO, BRAZIL
E-mail address: `mbel@impa.br`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FRIBOURG, PÉROLLES, CHEMIN DU MUSÉE 23, CH-1700 FRIBOURG, SWITZERLAND
E-mail address: `vincent.emery@gmail.com`